

SIZE 88MM X 220MM

Kloraxin Eye Ointment 1% w/w
(Chloramphenicol Eye Ointment BP 1% w/w)

Each gram contains:

Chloramphenicol B.P. 10mg

Description

White smooth ointment, free from foreign particles

Pharmacodynamics

Chloramphenicol is an antibiotic which is mainly bacteriostatic in action, but exerts a bactericidal effect against some strains of gram-positive cocci and against Haemophilus influenzae and Neisseria. It has a broad spectrum of action against both gram-positive and gram-negative bacteria, rickettsiae and Chlamydia.

Pharmacokinetics

Absorption:

Intraocular and some systemic absorption occur following topical application to the eye.

Distribution:

Aqueous humor- Measurable concentrations following topical application to the eye.

Indication

Indicated for the treatment of ocular infections involving the conjunctiva and/or cornea caused by the chloramphenicol susceptible organisms.

Chloramphenicol should be used only in serious infections for which less potent drugs are ineffective or contraindicated.

Recommended Dose

Usual adult and pediatric dose: To the conjunctiva, a thin strip (approximately 1cm) of ointment every three hours or more frequently.

Mode of Administration

Ophthalmic

Contraindication

Hypersensitivity

Warning and precautions

For ophthalmic use only. Use only as directed. Keep away from children. If irritation develops, or if redness is present, discontinue the use and consult a physician immediately. Do not swallow. If swallowed induce vomiting, call a physician. Prolonged or frequent intermittent use of Chloramphenicol topically should be avoided because of the possibility of hypersensitivity reactions, including bone-marrow hypoplasia. The content should be discarded 4 weeks after opening.

Interactions with Other Medicaments

By inhibiting microsomal enzymes, Chloramphenicol may interfere with the metabolism of dicumarol, phenytoin, tolbutamide and chlorpropamide.

Phenytoin and acetaminophen (paracetamol) may interfere with the metabolism of Chloramphenicol, causing toxic serum concentration. However, in a recent study, no interaction was noted with acetaminophen.

Phenobarbital and rifampin are known microsomal liver enzyme inducers, causing subtherapeutic concentrations of Chloramphenicol and, therefore, requiring dosage adjustments.

In vivo and in vitro studies of chloramphenicol used in conjunction with other antimicrobials reveal the possibility of antagonism: for example with β -lactams against Klebsiella pneumoniae, aminoglycosides against Escherichia coli, ampicillin against certain strains of Salmonella, penicillin against pneumococcus. This antagonism may be associated with the treatment failures and relapses.

Diuretics, such as mannitol, ethacrynic acid, and hydrochlorothiazide, increase the renal excretion of Chloramphenicol and its metabolites. These changes are clinically insignificant.

Pregnancy and Lactation

Pregnancy / Reproduction:

Problems in humans have not been documented.

Breast Feeding:

Problems in humans have not been documented

Side Effects

Hypersensitivity reactions, irritation

Symptoms and Treatments of Overdose

In the event of over dosage, no specific antidote is available; treatment is supportive and symptomatic.

If irritation develops, use of ointment should be discontinued and appropriate therapy instituted.

Storage Condition

Store below 30°C in a dry place. Keep the tube tightly closed.

Shelf Life

36 months from the date of manufacture.

4 weeks from the date of opening the tube.

Presentation:

Kloraxin 1% w/w Eye Ointment is available in a 5g Lacquered Aluminum collapsible tube.

5g x 25tubes is packed in outer shipper carton with package insert

Registration number: MAL09072753AZ

Revision date: 20.03.2020

Manufactured in India by:
Galentic Pharma (India) Pvt. Ltd
Plot No. 5-B & 5-C, Sector III,
Kandla SEZ, Gandhidham - 370230,
Gujarat.

Product Registration Holder and
Imported by: Unimed Sdn Bhd
No. 53 Jalan Tembaga SD5/2B,
Bandar Sri Damansara,
52200 Kuala Lumpur,
Malaysia.

KML1331PI-1