

PROSPAN®



Effervescent Cough Tablets

Information for the user

Prosparan® Effervescent Cough Tablets

Tablets

Active ingredient: Hedera Helix Extract (Dried Ivy Leaf Extract)

Read all of this leaflet carefully because it contains important information for you. This medicine is also available without prescription. However, you still need to use Prosparan® Effervescent Cough tablets as directed to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your doctor or pharmacist if you need more information or advice.
- You must contact a doctor if your symptoms worsen or do not improve after 7 days.
- If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

In this leaflet:

1. What Prosparan® Effervescent Cough Tablets are and what they are used for
2. Before you take Prosparan® Effervescent Cough Tablets
3. How to take Prosparan® Effervescent Cough Tablets
4. Possible side effects
5. How to store Prosparan® Effervescent Cough Tablets
6. Further information

1. What Prosparan® Effervescent Cough Tablets are and what they are used for

Prosparan® Effervescent Cough Tablets are traditionally used for relief of cough, throat irritation and reducing phlegm.

2. Before you take Prosparan® Effervescent Cough Tablets

Do not take Prosparan® Effervescent Cough Tablets

if you are allergic (hypersensitive) to the active substance or any of the other ingredients of Prosparan® Effervescent Cough Tablets.

Take special care with Prosparan® Effervescent Cough Tablets

This is a traditional medicine. Please consult your pharmacist/doctor if symptoms persist/worsen. Unsuitable for phenylketonurics. In the event of fructose intolerance, treatment should be carried out only after consultation with a doctor.

Taking other medicines

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines obtained without prescription.

Taking Prosparan® Effervescent Cough Tablets with food and drink

There are no special points to bear in mind. Can be taken with or without food.

Pregnancy and breast-feeding

Insufficient reliable data.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

No special precautions need to be taken into account.

Important information about some of the ingredients of Prosparan® Effervescent Cough Tablets

Each effervescent tablet contains 6.41 mmol (or 147.33 mg) sodium. Take this into account if you have to follow a low sodium diet. The aroma contains lactose. Each effervescent tablet contains 29.7 mg of lactose. Patients suffering from sugar intolerance should take Prosparan® Effervescent Cough Tablets only after consultation with a doctor.

3. How to take Prosparan® Effervescent Cough Tablets

Always take Prosparan® Effervescent Cough Tablets exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

Unless otherwise prescribed by your doctor, the usual dose is:

Adults and adolescents older than 12 years take 1 effervescent tablet 2 times daily. Children between 4-12 years take ½ effervescent tablet 3 times daily. The tablets are dissolved in a glass of hot or cold water (approx. 100-200 ml), and can be taken with or without food.

How long you should keep taking Prosparan® Effervescent Cough Tablets

Duration of treatment depends on the type and severity of the clinical picture. However, treatment should last for at least 1 week. Treatment with this medicine should be maintained for 2-3 days after symptoms have subsided to ensure lasting success of therapy.

If you have the impression that the effect of Prosparan® Effervescent Cough Tablets is too strong or too weak, talk to your doctor or pharmacist.

If you take more Prosparan® Effervescent Cough Tablets than you should

You should not exceed the daily dosage given in this leaflet or prescribed by your doctor. Taking considerably larger amounts (more than three times the daily dose) may cause nausea, vomiting or diarrhoea. In this case, you should consult your doctor.

If you forget to take Prospan® Effervescent Cough Tablets

Do not take a double dose to make up for a forgotten tablet. Simply continue to take Prospan® Effervescent Cough Tablets as prescribed by your doctor or as directed in this leaflet.

4. Possible side effects

Like all medicines, Prospan® Effervescent Cough Tablets can cause side effects, although not everybody gets them.

In very rare cases allergic reactions (shortness of breath, swelling, skin redness, itching) are known to occur after application of medicines containing ivy.

In sensitive persons gastric complaints may occur. If any of the side effects gets serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. How to store Prospan® Effervescent Cough Tablets

 **Keep out of reach of children.**
Jauhkan daripada capaian kanak-kanak.

Do not use Prospan® Effervescent Cough Tablets after the expiry date (EXP) which is stated on the carton and sachet. Store below 25°C. Protect from light and moisture. Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist on how to dispose of medicines no longer required. These measures will help to protect the environment.

6. Further information

What Prospan® Effervescent Cough Tablets contain

The active substance is heder helix extract (dried ivy leaf extract). Each 1.95g effervescent tablet contains 65 mg of heder helix extract from leaf (5-7.5:1). The other ingredients are anhydrous citric acid, sodium hydrogen carbonate, sodium carbonate, mannitol, simeticone, saccharin sodium 2 H₂O, acesulfame, aspartame, sodium citrate 2 H₂O, sorbitol, medium chain triglycerides, macroglycerol hydroxystearate, aroma. Each effervescent tablet contains 382 mg of carbohydrates (corresponding to 0.03 BU).

What Prospan® Effervescent Cough Tablets look like and contents of pack

Prospan® Effervescent Cough Tablets are brownish marbled effervescent tablets with a breaking notch. The effervescent tablets are single sealed in sachets and available in boxes of 6 and 20 tablets.

Manufacturer

Hermes Pharma GmbH
Hans-Urmiller-Ring 52, 82515 Wolfratshausen,
Germany.

Product Owner

Engelhard Arzneimittel GmbH & Co. KG
Herzbergstraße 3, 61138 Niederdorfelden,
Germany.

Product Registration Holder

Delfi Marketing Sdn. Bhd.
Level 6, Block A, Sky Park One City,
Jalan USJ 25/1, 47650 Subang Jaya,
Selangor, Malaysia.

This leaflet was last approved in May 2024

Note

An important term for describing the quality of extracts is the so-called “Drug to Extract Ratio” (DER). A DER of 5-7.5:1 means that 5 to 7.5 parts of the drug were needed to produce one part of extract.

We wish you a speedy recovery.

