

Amprol 12% w/v Solution

Composition: Each Amprol 12% 100ml solution contains amprolium hydrochloride 12 g (i.e. amprolium 10.6 g)

Product Description: Limpid clear-yellow solution

Pharmacodynamics/Pharmacokinetics : Amprolium hydrochloride is a coccidiostat belonging to the thiamine (Vitamin B1) analogue group that acts as a competitive antagonist of thiamine transport mechanisms. **ATC Vet Code: QP51AX09.** Therapeutic group: antiprotozoal.

Pharmacodynamics: Amprolium affects coccidia by competing with thiamine in their metabolic enzyme system, thus interfering with metabolism of carbohydrates necessary for multiplication or survival.

Pharmacokinetics: In the chicken, amprolium is rapidly absorbed following oral administration with C_{max} attained between 1 to 4 hours which is followed by rapid tissue distribution. Amprolium is eliminated rapidly mainly in the faeces with small amounts eliminated via the kidneys.

Indication: For the treatment and prevention of coccidiosis in poultry caused by *Eimeria spp.* Chickens (including broilers, breeders, replacement pullets and laying hens): *Eimeria tenella*, *E.necatrix*, *E.acervulina*, *E.praecox*. Moderate efficacy has been demonstrated against some strains of *Eimeria maxima*, *E.brunetti*, and *E.mitis*. Turkeys: *Eimeria adenoeides*, *E.gallopavonis* and *E.meleagrimitis*

Routes of administration: To be taken orally. Drinking water

	Dose (Amprol 12% Solution)	Dose (amprolium)	Duration of treatment
Initial treatment (Moderate infection)	50 ml / 50 litres of water	120 ppm	5-7 days
Initial treatment (Severe infection)	50 ml / 25 litres of water	240 ppm	5-7 days
Maintenance treatment (to follow initial treatment)	50 ml / 100 litres of water	60 ppm	1-2 weeks
Prophylaxis	50 ml / 100 litres of water	60 ppm	For duration of risk period

Contraindications: None.

Warnings and Precautions:

Special warnings for each target species: None.

Special precautions for use: None.

Special precautions for the disposal of unused product or waste material, if any – Any used or waste product material should be disposed of in accordance with national requirements.

Special precautions to be taken by the person administering the veterinary medicinal product to animals – In case of splashing into eyes, flush with plenty of water, seek for medical advice and show the package label where possible. Wash hands after use.

Interactions with other medicaments: No adverse events related to product interactions have been described.

Statement on usage during pregnancy and lactation: AMPROL 12% SOLUTION is safe to use in chickens and turkeys intended for breeding.

Adverse Effects: AMPROL 12% SOLUTION has a wide safety margin when used at the recommended dose rates.

Overdose and Treatment: In clinical and target species tolerance studies there have been no report of any adverse event associated with overdose. Antidote: thiamine

Storage Conditions: Do not store above 25°C.

Withdrawal Period: Chicken and Turkey: 0 days
Eggs: 0 days

Packing : 1 L, 5 L of plastic bottle.

FOR VETERINARY USE ONLY

Manufactured by :



DOPHARMA FRANCE

23 rue du Prieuré - Saint Herblon
44150 Vair sur Loire - FRANCE

Registration holder and repacker:



RHONE MA MALAYSIA SDN BHD (200001023279)

Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

Date of revision: December 2019
5LAMPRO3-01L 04/ XXX

LABEL CODE:	VERSION	COLORS
5LAMPRO3-01L		
PRODUCT: AMPROL 12%	Version: 4	Black
PACK SIZE: 1 L	MMM/YY: XXX	
COUNTRY: MY		
DIMENSIONS: cm = 10 x 14		! Please use the official PANTONE® (P) matching • system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE:	DATE + SIGNATURE: