

Vit C 50 Frx Tablet

DESCRIPTION

Oblong, Biconvex, Orange Tablet.

Size: 12.5 - 12.9mm (Length), 6.3 - 6.7mm (Width),
4.7-5.1mm (Thickness)

DOSAGE FORM

Tablets

COMPOSITION

Each tablet contains Ascorbic Acid 50mg

INDICATION

As a dietary supplement

RECOMMENDED USAGE

For Dietary Supplement

Children:

1 – 2 tablets per day (50 - 100 mg/day)

Adults:

2 – 10 tablets per day (100 - 500 mg/day)

Chew the tablet or allow to dissolve slowly in the mouth.

MODE OF ADMINISTRATION

Oral administration

CONTRAINDICATION

Hypersensitivity to the ingredients.

WARNING AND PRECAUTION

Patient with diabetes, prone to recurrent renal calculi (e.g. dialysis patients), undergoing stool occult blood tests and on sodium-restricted diets should not take excessive doses for extended periods of time. In large doses, ascorbic acid should be given with care to patients with hyperoxaluria.

Prolonged use of large doses of vitamin C may experience symptoms of deficiency of vitamin C when resume to normal intake. Excessive use of chewable vitamin C may lead to erosion of tooth enamel. Large dose of ascorbic acid have resulted in haemolysis in patients with G6PD deficiency.

SIDE EFFECT/ ADVERSE REACTION

Ascorbic acid is usually nontoxic and well tolerated. However, nausea, vomiting, heartburn, abdominal cramps, fatigue, flushing or redness of skin, headache, insomnia, a mild increase in urination, diarrhea, and sleepiness have been reported

DRUG INTERACTION

- Increased Effect: Ascorbic acid enhances iron absorption in the GI tract. Concomitant ascorbic acid taken with oral contraceptives-estrogen may increase contraceptive effect.
- Decreased Effect: Ascorbic acid and fluphenazine may decrease fluphenazine levels. Ascorbic acid and warfarin may decrease anticoagulant effect.
- Aspirin, nicotine, alcohol phenytoin, several appetite suppressants and some anticonvulsant drugs may decrease blood levels of ascorbic acid.

SIGNS AND SYMPTOM OF OVERDOSE AND TREATMENT

Gastrointestinal distress may indicate toxicity.

Forced fluids and diuresis may be beneficial in cases of toxicity due to massive ingestions of vitamin C.

STORAGE CONDITION

Store below 25°C and protect from light

Keep out of reach of children

Jauhi daripada kanak-kanak

SHELF LIFE

Please refer to the outer label.

PACKING/PACK SIZES

10 Aluminium bags of 100's in HDPE bottle.

Revision date: OCT 2009

MANUFACTURED BY:

The United Drug (1996) Co., Ltd., 208 Romklao Road, Minburi,
Bangkok 10510, Thailand.