

SELECTAN

300 mg/ml, solution for injection for cattle and swine

COMPOSITION PER ML:

Florfenicol	300 mg
Excipient q.s.ad	1 ml

A slightly yellowish and clear solution.



Florfenicol is a systemic broad spectrum antibiotic effective against most Gram-positive and Gram-negative bacteria isolated from domestic animals. Florfenicol acts by inhibiting protein synthesis at the ribosomal level and is bacteriostatic. However, in vitro studies of florfenicol demonstrate bactericidal activity against *Mannheimia haemolytica*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae*, and *Histophilus somni*.

In vitro testing has shown that florfenicol is active against the bacterial pathogens most commonly isolated in respiratory diseases in cattle (including *Pasteurella multocida*, *Mannheimia haemolytica*, *Histophilus somni* and *Arcanobacterium pyogenes*) and in pigs (including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*).

In cattle, intramuscular administration at the recommended dose of 20 mg/kg maintains efficacious blood levels for 48 hours. Maximum mean serum concentration (C_{max}) of 2.55 µg/ml occurs at 4.7 hours (T_{max}) after dosing. The mean serum concentration 24 hours after dosing was 1.4 µg/ml. The harmonic mean elimination half life was 26.2 hours.

In pigs, after initial intramuscular administration of florfenicol, maximum serum concentrations of between 1.9 and 3.1 µg/ml are reached after 2.2 hours and the concentrations deplete with a terminal mean half-life of 35.5 hours. After a second intramuscular administration, maximum serum concentrations of between 2.0 and 8.1 µg/ml are reached after 1.7 hours. Florfenicol concentrations achieved in lung tissue reflect plasma concentrations, with a lung: plasma concentration ratio of approximately 1. After administration to pigs by intramuscular route, florfenicol is rapidly excreted, primarily in urine. The florfenicol is extensively metabolised.

INDICATIONS:

Diseases caused by florfenicol susceptible bacteria:

Cattle: Therapeutic treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica*, *Pasteurella multocida*, and *Histophilus somni*.

Swine: Treatment of acute outbreaks of respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

DOSAGE AND ROUTE OF ADMINISTRATION:

Cattle: 20 mg/kg bodyweight (1 ml of SELECTAN per 15 kg) by intramuscular route to be administered twice 48 hours apart. The volume administered per injection site should not exceed 10 ml.

Swine: 15 mg/kg bodyweight (1 ml of SELECTAN per 20 kg) by intramuscular injection into the neck muscle twice at 48 hours intervals. The volume administered per injection site should not exceed 3 ml.

CONTRAINDICATIONS:

Do not use in adult bulls or boars intended for breeding purposes.

Do not use in cattle producing milk for human consumption.

Do not use in cases of hypersensitivity to the active substance or to any of the excipients.

INTERACTIONS:

None known.

PREGNANCY AND LACTATION:

Studies in laboratory species have not produced any evidence of teratogenic or foetotoxic effects. Safety during pregnancy and lactation has not been investigated in the target species. Use only accordingly to the benefit/risk assessment by the responsible veterinarian.

ADVERSE REACTIONS:

Cattle: A decrease in food consumption and transient softening of the faeces may occur during the treatment period. The treated animals recover quickly and completely upon termination of treatment. Administration of the product may cause inflammatory lesions at injection site which persist for 14 days.

Swine: Commonly observed adverse reaction are transient diarrhoea and/or peri-anal and rectal erythema/oedema which may affect 50% of the animals. These effects can be observed for one week. Transient swelling lasting up to 5 days may be observed at the site of injection. Inflammatory lesions at the injection site may be seen up to 28 days.

If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

OVERDOSE:

In swine, after overdosages, a reduction in feeding and hydration, weight gain and vomiting has been observed.

INCOMPATIBILITIES:

Do not mix the veterinary medicinal product with other veterinary medicinal products.

WITHDRAWAL PERIOD:

Cattle: Meat and offal: 30 days. Not permitted for use in lactating animals producing milk for human consumption.

Swine: Meat and offal: 18 days.

SPECIAL STORAGE PRECAUTIONS:

Keep out of the reach and sight of children.

Jauhi Ubat Dari Kanak-kanak.

Do not store above 25 °C.

Do not use after the expiry date stated on the label after Exp.

Shelf-life after first opening the container: 28 days.

SPECIAL WARNINGS(S):

The use of the veterinary medicinal product should be based on susceptibility testing and take into account official and local antimicrobial policies.

Do not use in piglets of less than 2 kg.

Swab the septum before removing each dose.

Use a dry, sterile syringe and needle.

Care should be taken to avoid accidental self-injection.

Avoid contact with eyes and skin.

If eye exposure occurs, flush eyes immediately with clean water.

If skin exposure occurs, wash the affected area with clean water.

Wash hands after use.

People with known hypersensitivity to florfenicol, propylene glycol or polyethylene glycol should avoid contact with the veterinary medicinal product.

DATE OF REVISION OF TEXT:

1st of January 2026

PRESENTATION:

Box with 1 glass bottle of 100 ml.

Box with 1 polypropylene bottle of 50 ml.

Box with 1 polypropylene bottle of 100 ml.

Box with 1 polypropylene bottle of 250 ml.

Registration holder:

HIPRA MALAYSIA SDN BHD

LOT NO. A-03-03A, LEVEL 3,

PLAZA MONT'KIARA, NO. 2,

JALAN KIARA, MONT'KIARA,

50480 KUALA LUMPUR,

WILAYAH PERSEKUTUAN KUALA LUMPUR

Reg. n° MAL10030014HA

FOR ANIMAL TREATMENT ONLY
To be supplied only on veterinary prescription
Controlled Medicine. Ubat Terkawal.

LABORATORIOS HIPRA, S.A.

Avda. la Selva, 135 . 17170 Amer (Girona) SPAIN

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com

HIPRA