

MEGLUXIN

Flunixin 50 mg/ml solution for injection

COMPOSITION PER ML:

Flunixin 50.000 mg

PHARMACODYNAMICS:

Flunixin meglumine is a non-steroid anti-inflammatory drug (NSAID) with analgesic and anti-pyretic activity. Flunixin meglumine acts as a non-selective and reversible inhibitor of cyclooxygenase (COX), an enzyme that converts arachidonic acid into unstable cyclic endoperoxides, which are then transformed into prostaglandins, prostacyclins and thromboxanes. Some of these prostanoids, such as the prostaglandins, participate in the physiopathological mechanisms of inflammation, pain and fever, which mean that their inhibition would be responsible for its therapeutic effects.

Due to the implication of prostaglandins in other physiological processes, the inhibition of the COX would also be responsible for different adverse reactions such as intestinal or renal damage. Prostaglandins form part of the complex processes involved in the development of endotoxic shock.

PHARMACOKINETICS:

Bovine: Flunixin meglumine, when administered intravenously to cattle, in a single dose of 2.2 mg/kg b.w., has a half-life of 4 hours. The maximum plasmatic level of Flunixin (14.9 µg/ml) is obtained in 10 minutes after intravenous administration of a dose of 2.2 mg/kg b.w. to calves, it then decreases to less than 0.1 µg/ml at 24 hours after being administered.

Flunixin meglumine shows a rapid distribution in highly irrigated tissues but equilibrium in tissues with less irrigation is established more slowly.

Flunixin initially undergoes a hydroxylation of the aromatic rings followed by conjugation. Afterwards, the conjugate may undergo an alkaline hydroxylation in the urine, thereby increasing the quantity of the free active ingredient.

Elimination is principally effected by the urinary route. An acid pH of the urine can increase the re-absorption of the drug in the renal tubules.

Equine: Flunixin meglumine, when administered by intravenous route to equids, in a single 1.1 mg/kg dose, has a plasmatic half-life of 2 hours.

Porcine: When administered to sows by intramuscular route at a dosage of 2.2 mg/kg b.w., the maximum plasmatic level is reached before an hour has elapsed and traces remain for 24 hours.

INDICATIONS:

Bovine: Indicated for the control of acute inflammation associated with respiratory disease. It has also been shown to have some benefit in the treatment of experimental acute bovine emphysema (Fog fever). It may be used as adjunctive therapy in the treatment of acute mastitis.

Equine: Indicated for the alleviation of inflammation and pain associated with musculo-skeletal disorders. It is also indicated for the alleviation of visceral pain associated with colic.

Porcine: Indicated for use as an adjunctive therapy in the treatment of swine respiratory diseases.

CONTRAINDICATIONS:

Do not use when signs of bloody discharges are present.

Do not use on animals that show hypersensitivity to flunixin meglumine.

Do not use on dehydrated, hypovolemic or hypotense animals.

SIDE EFFECTS:

None have been described at the dosages recommended.

Local reactions may occur locally after intramuscular administration.

TARGET SPECIES: Bovine, equine and porcine.

ADMINISTRATION ROUTE:

Intravenous in bovine and equine.

Intramuscular in swine.

DOSEAGE:

For use in equine musculo-skeletal disorders, the recommended dosage is 1 ml Megluxin per 45 kg bodyweight (equivalent to 1.1 mg flunixin per kg) injected intravenously once daily for up to five days according to clinical response.

For use in equine colic, the recommended dosage is 1 ml Megluxin per 45 kg bodyweight (equivalent to 1.1 mg flunixin per kg) by intravenous injection. Treatment may be repeated once or twice if colic recurs. The cause of colic should be determined and treated with concomitant therapy. For the treatment of endotoxaemia or septic shock associated with gastric torsion and with other conditions in which circulation of blood to the gastrointestinal tract is impaired, the recommended dosage is 1 ml per 200 kg bodyweight (0.25 mg/kg) administered every 6–8 hours.

For use in bovine acute inflammatory conditions, the recommended dosage is 2 ml Megluxin per 45 kg bodyweight (equivalent to 2.2 mg flunixin per kg) injected intravenously and repeated as necessary at 24 hour intervals for up to 5 consecutive days. The cause of the acute inflammatory condition should be determined and treated with concomitant therapy.

For use in swine, the recommended dosage is 2 ml Megluxin per 45 kg bodyweight (equivalent to 2.2 mg flunixin/kg) once by intramuscular injection, in the neck, in conjunction with appropriate antimicrobial therapy. The injection volume should be limited to a maximum of 5 ml per injection site.

SPECIAL PRECAUTIONS FOR USE IN ANIMALS:

Administer the product at room temperature (+15 to +25 °C).

Intravenous administration should be done very slowly.

Avoid intra-arterial administration to horses and cows. Horses accidentally injected intra-arterially may show adverse reactions such as ataxia, uncoordination, hyperventilation, excitation and muscular weakness. Water consumption during treatment and the animals' state of hydration should be watched during treatment, because there is greater risk of renal damage in cases of dehydration. Do not exceed the recommended dosage or duration of treatment. Use on animals younger than six weeks old or very old animals entails an additional risk. If it is not possible to avoid doing so, the animals should be given a reduced dose and given careful clinical follow-up. In intramuscular applications in swine, avoid depositing the product in the adipose tissue. It is preferable to not administer NSAIDs that inhibit the synthesis of prostaglandins to animals undergoing general anaesthesia until they are totally recovered.

WITHDRAWAL PERIOD:

Bovine: Meat: 7 days. Milk: 36 hours from last treatment.

Equine: Meat: 7 days from last treatment.

Porcine: Meat: 24 days from last treatment.

OVERDOSE:

May cause anorexia, diarrhoea, gastric ulcers, hypoproteinemia and renal necrosis. In such cases, suspend treatment and apply symptomatic therapy.

Signs such as lack of coordination and ataxia also appear.

INTERACTIONS:

Do not administer jointly, or with a difference of less than 24 hours, with other non-steroid anti-inflammatory drugs.

Some NSAIDs can bond in a large measure to plasmatic proteins and displace other drugs with affinity for bonding to them, which may cause toxic effects. This interaction is important for drugs with a narrow therapeutic margin: oral anticoagulants, methotrexate and some anticonvulsants such as phenytoin.

It may diminish the effect of some antihypertensives by inhibiting the synthesis of prostaglandins. Among them the diuretics, IECA, ARA and β-blockers should be pointed out.

Simultaneous administration with potentially nephrotoxic drugs should be avoided, especially cyclosporin.

It may diminish renal elimination of some drugs and increase their toxicity, as occurs with aminoglycosides.

SPECIAL PRECAUTIONS:

Store below 25 °C protected from light.

The product has a 28-day shelf life once the package has been opened.

Discard the used material and empty flasks by incineration.

Preserve the environment.

SPECIAL PRECAUTIONS TO BE TAKEN BY THE PERSON ADMINISTERING THE VETERINARY MEDICINAL PRODUCT TO ANIMALS:

To prevent possible reactions, avoid contact with skin. Gloves should be used while applying. Gloves should be used while applying. In case of contact with skin, wash immediately with water.

The product can cause reactions in those persons sensitive to it. In cases of known hypersensitivity to NSAIDs, do not handle the product. Reactions may be quite important.

Avoid the introduction of contamination while handling the product.

Acute pain and inflammation may occur in the case of accidental self-injection. Clean and disinfect the wound immediately, seek medical advice immediately and show the package insert or the label to the physician.

SPECIAL WARNINGS FOR EACH TARGET SPECIES:

The cause of the inflammation, pain or colic should be determined and treated concomitantly.

USE DURING PREGNANCY OR LACTATION:

Pregnancy: There are no specific studies on the target species, which means the risk/benefit analysis, should be assessed by the veterinarian before using it on gestating females.

DATE OF LAST TEXT REVISION:

1st of January 2026

PRESENTATIONS:

Vial with 50 ml.

Vial with 100 ml.

Vial with 250 ml.

Registration holder:

HIPRA MALAYSIA SDN BHD

LOT NO. A-03-03A, LEVEL 3,

PLAZA MONT'KIARA, NO. 2,

JALAN KIARA, MONT'KIARA,

50480 KUALA LUMPUR,

WILAYAH PERSEKUTUAN KUALA LUMPUR

Reg. n° MAL10030017HA

Keep medicine out of the reach of children.
Jauhi Ubat daripada Kanak-Kanak.
To be supplied only on veterinary prescription.
FOR VETERINARY USE ONLY

LABORATORIOS HIPRA, S.A.

Avda. la Selva, 135. 17170 Amer (Girona) SPAIN

Tel. (34) 972 43 06 60. Fax (34) 972 43 06 61. hipra@hipra.com

HIPRA