

For veterinary use only

Colmyc-E Inj 10%

Injectable solution



Composition
Enrofloxacin 10 g.
Excipient to 100 ml.

Pharmacodynamic properties

Enrofloxacin is bactericidal in action with activity against a wide range of Gram positive and Gram negative bacteria and mycoplasmas. The mechanism of action of the quinolones is unique among antimicrobials - they act primarily to inhibit bacterial DNA gyrase, an enzyme responsible for controlling the supercoiling of bacterial DNA during replication. Resealing of the double stranded helix is inhibited resulting in irreversible degradation of the chromosomal DNA. The fluoroquinolones also possess activity against bacteria in the stationary phase by an alteration of the permeability of the outer membrane phospholipid cell wall.

Molecular resistance to fluoroquinolones has been observed to arise from two principal sources, (i) alteration to DNA gyrase or topoisomerase IV and (ii) alterations in drug permeability of the bacterial cell. Both mechanisms lead to a reduced susceptibility of the bacteria to fluoroquinolones. Clinical resistance is dependent on several mutations accumulating in a step-wise manner.

Pharmacokinetic properties

The pharmacokinetics of enrofloxacin are such that oral and parenteral administration leads to similar serum levels. Enrofloxacin possesses a high distribution volume. Tissue levels 2-3 higher than that found in the serum, have been demonstrated in laboratory animals and target species. Organs in which high levels can be expected are the lungs, liver, kidney, skin, bone and lymphatic system. Enrofloxacin also distributes into the cerebrospinal fluid, the aqueous humour and the foetus in pregnant animals.

After an intravenous dose of 5 mg enrofloxacin per kg body weight (bw) to lactating dairy cattle, the total systemic exposure over the dosing interval of 24 h was at 7.1 mg*^h/L. In cattle serum, approximately 30% of drug exposure (2.31 mg*^h/L) consisted of ciprofloxacin, the active metabolite of enrofloxacin.

The drug was well distributed into the body compartments (V_{enro} = 1.5 L/kg, V_{cipro} = 8.51 L/kg). Total body clearance was 0.71 L/h/kg.

In milk, most of drug activity consisted of ciprofloxacin. Overall drug concentrations peaked at 4.1 mg/kg two hours after treatment. Overall drug exposure over 24 h was 22.1 mg*^h/L. The actives were eliminated from milk with a mean exposure half-life of 2.8 h.

Environmental properties

Not applicable.

Indications

Cattle

Diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. pasteurellosis, mycoplasmosis, coli-bacillosis, colisepticaemia and salmonellosis) and secondary bacterial infections subsequent to viral infections (e.g. viral pneumonia) where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.

Treatment of local signs (inflammation, milk quality and yield) associated with peracute/acute mastitis in lactating dairy cattle caused by *E. coli*, where herd history and previous sensitivity testing indicate enrofloxacin as the drug of choice.

Pigs

Diseases of the respiratory and alimentary tract of bacterial or mycoplasmal origin (e.g. pasteurellosis, mycoplasmosis, coli-bacillosis, colisepticaemia and salmonellosis) and multifactorial diseases such as atrophic rhinitis and enzootic pneumonia where clinical experience, supported where possible by sensitivity testing of the causal organism, indicates enrofloxacin as the drug of choice.

Target species

Cattle and pigs.

Recommended dosage

To ensure correct dosage, body weight should be determined as accurately as possible to avoid underdosing.

Cattle:

For respiratory and alimentary infections in cattle and secondary bacterial infections: administer by subcutaneous injection.

- 2.5 mg enrofloxacin per kg bodyweight daily by subcutaneous injection for 3 days (2.5 ml per 100 kg bodyweight). This rate may be doubled to 5 mg/kg bodyweight (5 ml per 100 kg) for 5 days for salmonellosis and complicated respiratory disease.

Not more than 10 ml should be administered at any one subcutaneous injection site.

For *E. coli* mastitis: administer by slow intravenous injection.

- 5.0 ml per 100 kg bodyweight (5 mg enrofloxacin per kg bodyweight) daily for 2 days.

Pigs:

For respiratory and alimentary infections in pigs and secondary bacterial infections: administer by intramuscular injection.

2.5 mg enrofloxacin per kg bodyweight daily by intramuscular injection for 3 days (2.5 ml per 100 kg bodyweight). This rate may be doubled to 5 mg/kg bodyweight (5 ml per 100 kg) for 5 days for salmonellosis and complicated respiratory disease.

Not more than 2.5 ml should be administered at any one intramuscular injection site in store pigs or 5 ml at any one intramuscular injection site in sows.

Route of administration

Subcutaneous or intravenous route.

Contraindications

- Do not administer in animals showing renal or hepatic insufficiencies, or in those sensitive to quinolones.
- Do not administer in newborn animals.

Side effects (frequency and severity)

- Rarely, alterations of the central nervous system may occur, as well as haematological and enzymatic disorders (transaminases).
- Articular lesions.

Special cautions for its use

Do not administer more than 10 ml. at the same site of injection.

Manufactured by:

s.p. veterinaria, s.a.

Ctra. Reus-Vinyols, km 4.1 • 43330 RIUDOMS (Spain)
Tel. +34 977 850 170 • Fax +34 977 850 405 • P.O.Box 60

Registration Holder:

LABORATORIOS REVEEX ASIA Sdn Bhd

Lot 3 & 5, Jalan Pjs 11/8

Bandar Sunway 46150 Petaling Jaya

Selangor Darul Ehsan - MALAYSIA

Phone: 60 3 563 33 493 - Fax: 60 3 563 23 410

Use during pregnancy and lactation

No contraindications were reported during these periods.

Interactions

- Antagonistic effects when administered with chloramphenicol, macrolides or tetracyclines can appear.
- Do not administer with nonsteroidal antiinflammatories.

Incompatibilities

Not reported.

Symptoms and treatment of overdosage

In case of overdosage, gastrointestinal and central nervous system disorders can appear. Discontinue the administration and carry out a symptomatic treatment.

Interactions with other medicinal products

- Antagonistic effects in the administration of chloramphenicol, macrolides or tetracycline can appear.
- Do not administer with nonsteroidal antiinflammatories.

Withdrawal period

Cattle: Subcutaneous Use.

Meat and offal: 10 days.

Milk: 84 hours (7 milkings).

Cattle: Intravenous Use.

Meat and offal: 4 days.

Milk: 72 hours (6 milkings).

Pigs: Intramuscular Use.

Meat and offal: 10 days.

Date of revision of package insert

23rd February 2010.

Proposed shelf life (as package for sale)

3 years.

Proposed shelf life (after first opening of the container)

28 days.

Proposed shelf life (after dilution)

Not applicable.

Storage conditions

Store in a cool, dry place (recommended below 25°C), protected from intensive light.

Presentations

100 ml. container.

Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Country Registration Nr.:

Keep out of the reach and sight of children

Controlled Medicine

Jauhi Ubat Dari Kanak - kanak

Distributed by:

F.E VENTURE Sdn BHD

Lot 3 & 5, Jalan Pjs 11/8

Bandar Sunway 46150 Petaling Jaya

Selangor Darul Ehsan - MALAYSIA

Phone: 60 3 563 33 493

Fax: 60 3 563 23 410

 **s.p. veterinaria, s.a.**