

CONTROLLED MEDICINE

KEEP OUT OF REACH OF CHILDREN

JAUHI UBAT DARI KANAK-KANAK

READ SAFETY DIRECTIONS

FOR ANIMAL TREATMENT ONLY

CYCOSTAT® 66G

ACTIVE CONSTITUENT: 66 g/kg Robenidine hydrochloride

PRODUCT DESCRIPTION: CYCOSTAT® 66G is a free flowing non dusty granule containing Robenidine hydrochloride. It is a safe and effective poultry feed additive which controls coccidiosis as well as fattens chickens.

PHARMACODYNAMICS/PHARMACOKINETICS:

The efficacy of CYCOSTAT® 66G is the result of a dual mode of action that is exerted upon different stages of the coccidial parasite as it develops in the intestinal mucosa of broiler chickens. Initially, robenidine hydrochloride is coccidiostatic; it arrests the development of the first generation schizonts. Secondly, robenidine hydrochloride is coccidiocidal, killing the second generation of schizonts and possibly the merozoites. Robenidine hydrochloride may also possess an effect on gametogony.

The absorption of robenidine in broiler chickens has been determined to be low, the total tissue residues were correspondingly found to be low. These factors, along with an essentially quantitative recovery of residue from tissue using an organic solvent, indicate that bound residues are negligible for robenidine.

INDICATION: For use in the control of coccidiosis in chickens caused by *Eimeria tenella*, *E. acervulina*, *E. necatrix*, *E. maxima*, *E. brunetti* and *E. mivati*.

RECOMMENDED DOSE: The period of administration is from day 1 to the start of the specified withdrawal period of 5 days prior to slaughter.

Thoroughly mix CYCOSTAT® 66G into the feed at the rate of 500g per tonne to give a concentration of 33 g/t Robenidine in the complete feed.

MODE OF ADMINISTRATION: Oral - Use continuously in the feed.

CONTRAINDICATION:

DO NOT use in birds laying eggs for human consumption.

DO NOT use in feeds containing Bentonite.

WARNINGS AND PRECAUTIONS:

Feed continuously as sole ration.

Avoid inhaling dust during mixing. Avoid contact of product with skin, eyes and clothing. Wash thoroughly after handling.

INTERACTIONS WITH OTHER MEDICAMENTS: There is no known evidence of any incompatibilities or interactions with other products.

PREGNANCY AND LACTATION: Not Applicable.

UNDESIRABLE EFFECTS: None known.

OVERDOSE AND TREATMENT: If an overdose is suspected seek advice from a veterinarian.

STORAGE: Store below 25°C away from sunlight. The bag should be closed securely after use.

DOSAGE FORM: CYCOSTAT® 66G is an oral premix for inclusion in the feed.

PACKAGING: Each lot is packaged in 20 kg multiwalled paper bags.

DISPOSAL OF CONTAINER: Shake and empty contents into medicated feed. Shred and bury empty containers in a local authority landfill. Empty containers and product should not be burnt.

WITHHOLDING PERIOD: Remove all medicated feed 5 days before slaughter for human consumption

MANUFACTURED BY

Zoetis Medolla Manufacturing S.r.l.

Via Rubadello No. 6

I-41036 Medolla

ITALY

MARKETING AUTHORISATION HOLDER:

Pahang Pharmacy Sdn. Bhd

Lot 5979, Jalan Teratai

5 1/2 Mile off Jalan Meru, 41050 Klang Selangor

MALAYSIA

Date of Revision: 27.04.2022 Reg. No.: MAL10070030HA

AHZ 979 1310