

Package Insert

1. Product name

Pizogran 0,5 Tablets

2. Name and Strength of Active Ingredient

Pizotifen 0,5 mg (as Malate)

3. Product description

Film-Coated Tablets.

White, round, film-coated tablets with Remedica's logo on one side.

4. Pharmacological Properties

Pharmacodynamics

Pharmacodynamic studies demonstrate pizotifen to have powerful anti-serotonin and anti-tryptaminic properties, marked anti-histaminic effects and some antagonistic activity against kinins. It also possesses weak anti-cholinergic effects and sedative properties.

Pizotifen also possesses appetite-stimulating properties.

The prophylactic effect of pizotifen in migraine is associated with its ability to modify the humoral mechanisms of headache.

It inhibits the permeability-increasing effect of serotonin and histamine on the affected cranial vessels, thereby checking the transudation of plasmakinin so that the pain threshold of the receptors is maintained at 'normal' levels. In the sequence of events leading to migraine attack, depletion of plasma serotonin contributes to loss of tone in the extracranial vessels. Pizotifen inhibits serotonin re-uptake by the platelets, thus maintaining plasma serotonin and preventing the loss of tone and passive distension of the extracranial arteries.

Pharmacokinetics

Absorption

Absorption of pizotifen is fast (absorption half life 0.5 to 0.8 hours) and nearly complete (80%).

Biotransformation

Pizotifen is metabolised with a half life of about 1 hour. The main metabolite (N-glucuronide) is eliminated with a half life of approximately 23 hours.

Distribution

Protein binding amounts to 91% and distribution volume to 485 litres.

Elimination

Less than 1% of the administered dose is excreted unchanged in the urine, whereas 55% is excreted as metabolites

5. Clinical Particulars

Indications

Prophylactic (interval) treatment of vascular headaches, such as – typical or atypical migraine – vasomotor headaches - cluster headaches (Horton syndrome). It is not effective in relieving migraine attacks once in progress.

Posology and method of administration

Adults

Starting with 0,5mg daily, the dosage should be progressively increased. The average maintenance dosage is 1,5mg daily in divided doses as a single dose at night. In refractory cases, the physician may gradually raise the dosage to 3-4,5mg daily taken in 3 divided doses.

Method of administration

Oral.

Contraindications

Known hypersensitivity to pizotifen or any of the excipients

Warnings and Precautions

Although the anticholinergic activity of pizotifen is relatively weak, caution is required in the presence of closed angle glaucoma and in patients with a predisposition to urinary retention. Dosage adjustment may be necessary in patients with kidney insufficiency. Patients with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Interactions with other medicaments

The central effects of sedatives, hypnotics, antihistamines (including certain common cold preparations) and alcohol may be enhanced by pizotifen.

Pregnancy and Lactation

As clinical data with pizotifen in pregnancy are very limited it should only be administered during pregnancy under compelling circumstances.

Although the concentrations of pizotifen measured in the milk of treated mothers are not likely to affect the infant, its use in nursing mothers is not recommended.

Effects on ability to drive and use machines

Patients should be cautioned about the possibility of drowsiness and informed of its significance in the driving of vehicles and the operation of machinery.

Undesirable effects

The most commonly occurring side-effects are drowsiness and an increased appetite which may lead to an increase in body weight. Other side-effects such as dizziness, dry mouth, nausea, constipation, hallucinations and paraesthesia have been reported infrequently. Rare instances of sleep disorders, depression and other mood disturbances such as anxiety and aggression have occurred. In children CNS stimulation may occur. Rare cases of hypersensitivity reaction such as rash, facial oedema and urticaria have been reported.

Symptoms and treatment of overdose

Symptoms: Drowsiness, nausea, hypotension, dizziness, excitatory states (in children), respiratory depression, convulsions (particularly in children), coma.

Treatment: Administration of activated charcoal is recommended; in case of very recent intake, gastric lavage may be considered. If necessary, symptomatic treatment including monitoring of the cardiovascular and respiratory systems. For excitatory states or convulsions, benzodiazepines may be used.

6. Pharmaceutical Particulars

Storage conditions

Store below 30°C. Protect from light and moisture.

Shelf-life

36 months

Pack size available

100 tablets in PP/PE containers

60 (6 x 10), [100 (10 x 10)-Malaysia] and 1000 (10 x 100) tablets in Blister packs

Not all presentations are available locally.

7. Manufacturer

Remedica Ltd.
Aharnon Street, Limassol Industrial Estate,
3056 Limassol, Cyprus.

8. Date of revision of the text

07/2022

Product Registration Holder

Eucogen Sdn. Bhd.,
No. 6A, Jalan Sungai Burung U 32/U, Bukit Rimau, Seksyen 32, 40460 Shah Alam.
Telephone Number: 03 5131 1997
Fax Number: 03 5120 1639

For internal use only: my-pi-pizogran-0.5mg-fc-tabs-a2