

**FLEXIT SR****Glucosamine Hydrochloride
Sustained Release Tablets****PRESENTATION:**

Sustained release Tablet

QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film coated tablet contains Glucosamine Hydrochloride USP 1500 mg

PHARMACEUTICAL FORM

Tablet

CLINICAL PARTICULARS**Therapeutic indication:**

Adjuvant therapy for osteoarthritis.

Dosage and Method of Administration

Adults: One FLEXIT SR Tablet once daily with food and with a glass of water, or as prescribed.

Duration of treatment: 3 months.

The treatment course may be repeated at intervals of 2 months.

Method and route of administration: Oral route.

Contra-indications:

FLEXIT SR is contraindicated in patients with hypersensitivity to any of its ingredients, during pregnancy and lactation.

Special Warnings and Special Precautions for Use

Use of Glucosamine must be with caution if there is history of allergy to shellfish and shellfish products. Asthma patients may be at risk for an asthma exacerbation when taking the combination of glucosamine. Diabetics should use Glucosamine cautiously as glucosamine may affect insulin sensitivity or glucose tolerance.

Glucosamine must be taken with caution in those taking oral hypoglycemics, doxorubicin, etoposide and teniposide. The usefulness of glucosamine in children has not been established.

It is advisable to take an anti-inflammatory drug in the initial treatment of osteoarthritis together with glucosamine.

Interaction with Other Medicaments and Other Forms of Interaction

Glucosamine when co-administered can decrease the efficacy of oral hypoglycemics, doxorubicin, etoposide and teniposide.

Undesirable Effects**Cardiovascular**

Peripheral oedema, tachycardia were reported in a few patients following larger clinical trials investigating oral administration in osteoarthritis.

Causal relationship has not been established.

Central Nervous System :

Drowsiness, headache, insomnia have been observed rarely during therapy (less than 1%)

Gastrointestinal :

Nausea, vomiting, diarrhea, dyspepsia or epigastric pain, constipation, heartburn and anorexia have been described rarely during oral therapy with glucosamine



**Skin :**

Skin reactions such as erythema and pruritus have been reported with therapeutic administration of glucosamine.

Pregnancy and Lactation

In pregnancy and lactation, seek medical advice before use.

Overdosage:

No cases of accidental or intentional overdose are known. The animal acute and chronic toxicological studies indicate that toxic effects and symptoms toxicity are not likely to occur, even after high overdoses

PHARMACOLOGICAL PROPERTIES**Pharmacodynamic Properties**

Glucosamine stimulates biosynthesis of cartilage and joint substance so as to benefit osteoarthritis. Its intake results in a dose-dependent increase in proteoglycan as well as collagen production by chondrocytes. Glucosamine also stimulates hyaluronic acid synthesis, and thereby increases viscosity of synovial fluid. Besides, glucosamine diminishes cartilage degeneration by reducing the production of cartilage-damaging superoxide radicals and destructive enzymes.

Taken for prolonged periods, Glucosamine relieves the pain and inflammation associated with osteoarthritis by regenerating the cartilage matrix and increasing its and synovial fluid's hyaluronic acid content

Pharmacokinetic Properties

Glucosamine when taken orally, ionizes completely within the stomach by HCl to form glucosamine ions. It is 90% absorbed from the small intestine. However, it undergoes extensive first-pass metabolism in gut and/or liver resulting in approximately 26% bioavailability. The plasma half-life of glucosamine is short, less than an hour.

Glucosamine has a special affinity for cartilage and is incorporated by the chondrocyte into the extracellular matrix in about 4 hours. Approximately 8-12% of the absorbed glucosamine is retained in tissues such as cartilage. Glucosamine is also concentrated in liver and kidneys.

2-10% of the administered glucosamine is lost in feces between 24 to 72 hours. Of the absorbed glucosamine, 20-30% subsequently appears in urine, up to 70% as exhaled carbon dioxide. The maximum excretion achieved is between 4 and 8 hours. The elimination half-life of glucosamine is 28 hours.

Shelf-Life

The expiry date is indicated on the packaging.

Special Precautions for Storage

Store below 30°C, away from direct sunlight.

Nature and Contents of Container

30 Tablets packed in 185 ml OFC white round screw type HDPE containers.

Further information available upon request.

KEEP OUT OF REACH OF CHILDREN

Package insert revised in July 2025

Flexit SR is a trademark of:



MEDREICH PLC
Warwick House, Plane Tree Crescent,
Feltham TW13 7HF, UK

Product Registration Holder:
Healol Pharmaceuticals Sdn. Bhd.
74-3, Jalan Wangsa Delima 6,
KLSC Wangsa Maju,
53300 Kuala Lumpur, Malaysia.

Mfd. by: ADCOCK INGRAM LIMITED
(A Medreich Group Company)
49 C And D Industrial Area, Anekal Taluk,
Bommasandra, 560 099, INDIA.



ADCOCK INGRAM LIMITED

CUSTOMER: **Healol Pharmaceuticals Malaysia**

COMPONENT: **Flexit 1500mg SR Tablets S/L English PIL- WA**

ITEM CODE: **1229708** (AWR/Q1/ME/25/0001)

DIMENSION: **110 x 190 mm (±2mm)**

ARTWORK SCALE: **100%**

REASON FOR CHANGE: **Change in text matter and dimensions tolerance updated.**

COLOUR SCHEME: **2 - Pantone Red 032 C and Black**

SUPERSEDED ITEM CODE: **1216333**

SPECIFICATION: **Printed on 40 to 45 GSM news print paper**

Head - PDC

Head - TT

Head - QC

Head- Production

Head CQA/QA

Prepared by:
Date:

Checked by:
Date:

Checked by:
Date:

Approved by:
Date:

Authorized by:
Date: