

SYMPTOMS AND TREATMENT FOR OVERDOSE AND ANTIDOTE(S):

Overdosage is unlikely, if accidentally ingested, gastric lavage should be performed.

PACKING / PACK SIZES:

Plastic bottle of 100 ml, 120ml and 200ml

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS:

Store below 30°C in a tight, light-resistant container. Protect from light.

SHELF LIFE: 3 years

NAME AND ADDRESS OF MANUFACTURER:

SM PHARMACEUTICALS SDN BHD (218620-M)

Lot 88, Sungai Petani Industrial Estate

08000 Sungai Petani

Kedah Darul Aman,

Malaysia.

17.05.2023

SM PHARMACEUTICALS SDN BHD

CROTOCORT LOTION

(CROTAMITON 100 MG)

DESCRIPTION:

A white, slightly viscous, homogenous lotion

COMPOSITION:

Each ml contains – Crothamiton 100 mg and Benzyl Alcohol 0.1% as preservative.

ACTIONS AND MODE OR MECHANISMS OF ACTION:

Antipruritic and acaricide.

Crothamiton is effective against various forms of pruritus. The relief it affords sets in rapidly and lasts for some six hours. By relieving the itching, it prevents irritation of the skin caused by scratching and thus reduces the risk of secondary infections. Since crothamiton is also an acaricide and, in addition, inhibits the growth of streptococci and staphylococci, it is particularly suitable for the treatment of scabies, including secondary pyodermal infections.

PHARMACOLOGY:

Crothamiton penetrates easily into the skin without leaving the greasy or shiny deposit on the surface.

The lotion is preferable for application to the hairy and exposed parts of the body and for use on weeping skin.

INDICATIONS:

- a) Symptomatic relief of pruritus (itching) of varying origin, e.g., essential pruritus, senile pruritus, anogenital pruritus, pruritus associated with allergies, jaundice or diabetes mellitus and pruritus due to insect bites and stings.
- b) Scabies (infestations of the skin by mites).

CONTRAINDICATIONS:

Hypersensitivity to the active substance or excipients.

SIDE EFFECTS / ADVERSE REACTIONS:

Crotamiton is generally well-tolerated. Only in exceptional cases is irritation of the skin or contact allergic liable to occur; in such instances the preparation should be withdrawn.

PRECAUTIONS / WARNINGS:

Crotamiton should not be allowed to come into contact with the conjunctiva. Where such contact occurs by accident, the eye should be bathed with water. Treatment with the cream is contraindicated in acutely inflamed, weeping skin conditions or skin wounds owing to the danger of systemic absorption.

Pregnancy / Reproduction:

Studies have not been done in either animals or humans; however risk-benefit must be considered (FDA Pregnancy Category C)

Breast-feeding:

Problems in humans have not been documented; however risk benefit must be considered.

Use in pregnancy and lactation

Since the risk-benefit ratio in pregnancy is inadequately documented, use of crotamiton during pregnancy, particularly during the first trimester is not advised. It is not known whether the active substance passes into the breast milk. Nursing mothers should, in all events, avoid applying crotamiton in the area of the nipple.

DIRECTIONS FOR USE:

i)

Pruritus:

Crotamiton lotion should be rubbed into the affected parts 2 – 3 times a day until itching has subsided. This also applies to the refractory cases requiring more prolonged treatment, as crotamiton is well tolerated by the skin. Patients using crotamiton as self-medication should consult a doctor if the itching still persists after 5 days treatment.

ii)

Scabies:

After the patient has first taken a warm bath, the skin should be well dried and crotamiton then rubbed into the entire body surface (excluding the face and scalp) until no traces of the preparation remain visible on the surface. The application should be repeated once daily, preferably in the evening, for total of 3 – 5 days depending on the response. When applying crotamiton, special attention should be paid to sites that are particularly susceptible to infestation by the mites (interdigital space, wrists, axillae and genitalia). Areas where there is pus formation should be covered with a dressing impregnated with crotamiton. While the treatment is in progress the patient may take a bath shortly before the next application.

After completion of the treatment, a cleansing bath should be taken, followed by a change of bed linen and under clothing. In small children, crotamiton should never be applied more than once a day.