

CLON POWDER 100mg/g

Ingredient(s):

Each g contains :
Tiamulin Hydrogen Fumarate 100mg

Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):

Tiamulin is usually a bacteriostatic antibiotic, but can be bactericidal in very high concentrations (more than 100 times the MIC) against susceptible organisms. The drug acts by binding to the 50S ribosomal subunit, thereby inhibiting bacterial protein synthesis.

Tiamulin has good activity against many gram-positive cocci, including most *Staphylococci* and *Streptococci* (not group D streps). It also has good activity against *Mycoplasma* and spirochetes. With the exception of *Haemophilus spp* and some *E.coli* and *Klebsiella* strains, the drug's activity is quite poor against gram-negative organism.

Tiamulin is well absorbed orally by swine. Approximately 85% of a dose is absorbed and peak levels occur between 2~4 hours after a single oral dose. In rats, dogs, and probably also by chickens, tiamulin was absorbed to the extent of more than 90% after oral application. Tiamulin is apparently well distributed, with highest levels found in the lungs. Tiamulin is extensively metabolized to over 20 metabolites, some having antibacterial activity. Approximately 30% of these metabolites are excreted in the urine with the remainder excreted in the feces. More than 95% of the dose was excreted by all animal species with a half-life of less than 1 day. In pigs or chickens, less than 1% of the dose was excreted with a half-life of 3~4 or 6~8 days, respectively. In rats, dogs, chickens and very likely pigs, two-third of the dose reached the feces through the bile.

Indication(s):

Swine:

For the control of diseases such as Swine dysentery (SD) caused by *Treponema hyodysenteriae*, Swine enzootic pneumonia (SEP) caused by *Pasteurella multocida* and *Mycoplasma hyopneumoniae*, and pleuropneumonia caused by *Actinobacillus pleuropneumoniae*.

Poultry:

For the control of Chronic respiratory disease (CRD), Infectious synovitis and sinusitis.

Target Species

Swine and non-laying chicken.

Dosage and Administration:

Control of Swine dysentery:

Prevention: 0.3~0.6kg/ton feed, use for 30 consecutive days.

Treatment: 1kg/ton feed, use for 7 consecutive days.

Control of Swine enzootic pneumonia (SEP):

Prevention: 0.5~1.0kg/ton feed, use for 30 consecutive days.

Treatment: 2kg/ton feed, use for 10 consecutive days.

Control of pleuropneumonia:

Prevention: 0.3~0.5kg/ton feed, use for 30 consecutive days. (may in combination with 100~150ppm CTC or OTC)

Treatment: 1kg/ton feed, use for 10 consecutive days. (may in combination with 300ppm CTC or OTC)

Control of Chronic respiratory disease (CRD), Infectious synovitis and sinusitis:

Prevention: 0.5kg/ton feed for 7 days every month, or 0.2kg/ton feed continuously.

Treatment (non-laying chickens): 2kg/ton feed for 5 days.

Orally via feed mixing.

Contraindications:

Tiamulin should not be administered simultaneously with polyether ionophores (e.g., monensin, lasalocid, narasin or salinomycin) at level above 30ppm as signs of interaction may appear.

Precaution(s) / Warning(s):

1. Not for use in swine over 250 lbs of body weight.
2. Do not use in birds which are producing eggs for human consumption.

Interaction with Other Medicaments

- 1) Tiamulin should not be administered to animals having access to feed containing polyether ionophores (e.g., monensin, lasalocid, narasin, or salinomycin) as adverse reactions (growth retardation, ataxia, paresis, and paralysis of the legs, and in the case of young chickens, an increases mortality rate) may occur.
- 2) Concomitant use with other antibiotics that bind to the 50S ribosome (e.g., clindamycin, lincomycin, erythromycin, tylosin) could lead to decreased efficacy due to competition at the site of action.
- 3) The following coccidiostats may be administered concurrently with tiamulin: zoalen, amprolium, clopidol, robenidin, buquinolat, decoquinat, and furazolidine.
- 4) No signs of incompatibility were observed when tiamulin was administered concurrently with the following therapeutic agents and additives in their usual dosage: chlortetracycline, neomycin, oxytetracycline, nitrovin, carbodox, dimetridazole, sulfadimidine, sulfathiazole sodium, and furazolidone.

Pregnancy and Lactation

Teratogenicity studies done in rodents demonstrated no teratogenic effects at doses up to 300mg/kg. The drug is not tumorigenic, carcinogenic, teratogenic or mutagenic.

Side Effect(s) / Adverse Reaction(s):

- 1) Adverse effects occurring with this drug at usual doses are considered unlikely.
- 2) Rarely, redness of the skin, primarily over the ham and underline, has been observed.
- 3) Adverse reactions of growth retardation, ataxia, paresis, and paralysis of the legs, and in the case of young chickens, an increases mortality rate may occur when tiamulin was administered to animals having access to feed containing polyether ionophores (e.g., monensin, lasalocid, narasin, or salinomycin).

Environmental Property

None has been mentioned.

Symptoms and Treatment for Overdosage, and Antidote(s):

Oral overdoses in pigs may cause transient salivation, vomiting and CNS depression (calming effect). Discontinue drug and treat symptomatically and supportively if necessary.

Shelf-Life:

4 years.

Storage Condition(s):

Store at temperature below 30°C.

Product Description and Packing(s):

A white, fine powder and without other foreign matter.

Drum of 10Kg

To amend from
Drum of 10kg to "
10kg in Paper bag".

For food producing animals product:

Withdrawal Periods

Poultry: 7 days before slaughter.

Swine: 5 days before slaughter.

Do not use in birds which are producing eggs for human consumption.

Maximum Residual Limit (MRL):

Porcine: Muscle 100µg/kg, liver 500µg/kg.

Chicken: Muscle 100µg/kg, skin+fat 100µg/kg, liver 1000µg/kg, egg 1000µg/kg.

Turkey: Muscle 100µg/kg, skin+fat 100µg/kg, liver 300µg/kg.



Made by YSP

Manufacturer:

Y.S.P. INDUSTRIES VIETNAM CO., LTD

Road No. 6, Nhon Trach 3 Industrial Zone, Nhon Trach,
Dong Nai Province, Vietnam Tel: +8461 3569 993

Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (192593 U)

Lot 3, 5 & 7, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor Darul Ehsan, Malaysia.

Ordering Line: 1 800 88 3027 Product Info: 1 800 88 3679

Date of revision: 21 January 2016

Revise the date
of revision.