

TYLONE 668MG Water Soluble Powder

FOR ANIMAL USE ONLY

Product Name :

TYLONE 668MG Water Soluble Powder

Name and Strength of Active Substance

Each gram of TYLONE 668MG water soluble powder contains Tylosin Tartrate 668mg

Product Description

TYLONE 668MG Water Soluble Powder is a white to slightly yellowish powder with characteristic odour and bitter taste, containing 66.8% w/w Tylosin tartrate.

Pharmacodynamics

Tylosin is a macrolide antibiotic produced by a strain of *Streptomyces fradiae*. It exerts its antimicrobial effect by inhibiting protein synthesis of susceptible micro-organisms. The tylosin spectrum of activity includes Gram-positive bacteria including *Clostridium perfringens* and some Gram-negative strains such as *Pasteurella* and *Mycoplasma* spp. at concentrations of 16µg/ml or less.

Pharmacokinetics

Absorption: Tylosin reaches maximal blood levels between 1 and 3 hours after an oral dose. Minimal or no blood levels remain 24 hours after an oral dose.

Distribution: After oral doses were given to pigs, tylosin was found in all tissues, between 30 minutes and two hours after administration, except for the brain and spinal cord. Biotransformation and Elimination: It has been shown that most of the material which is excreted is to be found in the faeces and consists of tylosin (factor A), relomycin (factor D) and dihydrodesmycosin.

Indication

For the treatment of chronic respiratory disease (CRD) associated with *Mycoplasma gallisepticum* sensitive to tylosin in broiler and replacement chickens. For the control of CRD associated with *Mycoplasma gallisepticum* sensitive to tylosin at the time of vaccination or other stress in chickens. For the control of CRD associated with *Mycoplasma synoviae* sensitive to tylosin in broiler chickens.

Recommended Dosage

Poultry : Dissolve 150 g into 160 Litres or 500 g into 534 litres of drinking water.

Administer from 1 to 5 days as sole source of drinking water. Treated chickens should consume enough medicated drinking water to provide 50 milligrams of tylosin per pound of body weight per day.

Mode of Administration

To be administered via animal drinking water. Mixing Directions : To assure thorough dissolution, place the TYLONE 668MG powder in a small mixing container and add 25 Litres of water . Stir until all the powder completely dissolved. When mix the product with water, always add water to the powder. Do not pour the powder into the water. Mix this concentrated stock solution with water to make the final desired volume of medicated animal drinking water. Prepare fresh medicated solution daily.

Contraindications

Tylosin is contraindicated in patients hypersensitive to it or other macrolide antibiotics (e.g., erythromycin). Tylosin is contraindicated in horses, as severe and sometimes fatal diarrheas may result from its use in that species.

Warnings and Precautions

Tylosin is contraindicated in patients hypersensitive to it or other macrolide antibiotics (e.g., erythromycin). Not to be handled by person with known macrolide antibiotic allergy. Use appropriate personal protection. Wear impervious gloves when handling powder. Open sachet with care and not in windy area to avoid forming dust cloud. Use adequate masks and safety goggles in case of powder being blown.

Respiratory Protection	:	Wear dust mask in case powder being blown.
Hands	:	Wear Impervious gloves when handling powder.
Eye Protection	:	Wear Safety goggles when handling powder.
Skin Protection	:	Wear protective clothing to minimize skin exposure.

Drug Interactions

Drug interactions with tylosin have not been well documented.

Statement on usage during pregnancy and lactation

No information was located with regards to the reproductive safety of tylosin, but it is unlikely to have serious teratogenic potential. No information was located with regards to the safety use of tylosin during lactation. Do not use in layers producing eggs for human consumption.

Adverse Effects / Undesirable Effects

Most likely adverse effects with tylosin are pain and local reactions at intramuscular injection sites, and mild GI upset (anorexia, and diarrhea). Tylosin may induce severe diarrheas if administered orally to ruminants or by any route to horses. In swine, adverse effects reported include edema of rectal mucosa and mild anal protrusion with pruritis, erythema, and diarrhea.

Overdose and Treatment

Tylosin is relatively safe in most overdose situations. The LD50 in pigs is greater than 5 g/kg orally, and approximately 1 g/kg IM. Dogs are reported to tolerate oral doses of 800 mg/kg. Long-term (2 year) oral administration of up to 400 mg/kg produced no organ toxicity in dogs. Shock and death have been reported in baby pigs overdosed with tylosin.

Withdrawal Period(s)

5 days

Shelf-life

3 years from the date of production.

Allowable Maximum Residual Limit (MRL)

All food-producing species : Milk (cattle) = 0.05mg/kg, Muscle, liver, kidney (chicken and cattle), edible tissue (cattle), fat (chicken), egg (chicken) = 0.20mg/kg [Ref. REGOVP, Version 2, dated Aug 2008]

Storage Conditions

Store in a dry place below 25°C. Keep out of reach of children. Jauhkan dari kanak-kanak. Any opened sachet or unused medicated water to be discarded after 24 hours.

Dosage Forms and packaging available

Available in 150 g, 500 g, 1 kg powder per sachet, pack in 10, 20, 50 or 100 sachets per box.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2010-08-16