

# CHEMIX POWDER 500mg/g

**Ingredient(s):**

Each g contains :

Flumequine ..... 500mg

**Pharmacology (Summary of Pharmacodynamics and Pharmacokinetics):**

Flumequine is a synthetic antibiotic belong to fluoroquinolone group. It is active against Gram negative bacteria and is mainly used for treatment of enteric infections in domestic animals. After oral administration, it quickly passes into the bloodstream and is widely dispersed in the tissues with a high affinity for lung & kidney. It is mainly eliminated through urine and feces.

After oral administration of 12mg/kg, the plasma level after the first administration reached a plateau at 0.5 hours post dosing (2.5-5.8mg equiv/L) and then fell to 0.8-1.5mg equiv/L immediately prior to the next dose 12 hours later. Between 52-73% of administered radioactivity was eliminated in the urine (49-70% in the first 24 hours) while faecal elimination accounted for a further 21-36%. Total recovery of radioactivity amounted 52-73% at 168 hour post dosing.

Thus, Flumequine is rapidly absorbed after oral administration and most rapidly eliminated within 168 hours post-dosing.

**Indication(s):**

For the prevention and treatment of diseases in poultry, swine, goat, sheep and calf such as enteric and septicaemic diseases due to *E.coli*, *Salmonella*; respiratory diseases due to *Bordetella*, *Haemophilus*, *Pasteurella*.

**Dosage and Administration:**

Prevention: 100mg/ L of water (50ppm)

Treatment: 200mg/L of water (100ppm)

Oral administration via drinking water for 3-5 days.

**Contraindications:**

Severe liver or kidney insufficiency.

**Precaution(s) / Warning(s):**

Not to be used in animals from which eggs are produced for human consumption

**Interaction with Other Medicaments:**

Nitrofurantoin may impair the efficiency of flumequine. Antagonistic effect with trimethoprim, sulfonamides, furazolidone, phenylbutazone & hydrocortisone.

**Pregnancy and Lactation:**

None has been mentioned.

**Side Effect(s) / Adverse Reaction(s):**

Rarely hypersensitivity to Flumequine may occur.

**Environmental Property:**

Non-hazardous

**Symptoms and Treatment for Overdosage, and Antidote(s):**

None has been mentioned.

**Shelf-Life:**

3 years.

**Storage Condition(s):**

Store at temperature below 30°C.

**Product Description and Packing(s):**

A white to off-white color powder, and without other foreign matter.

Aluminium Pouch of 100g.

**For food producing animals product:****Withdrawal Periods**

7 days

**Maximum Residual Limit (MRL):**

Bovine, Ovine, Porcine:

Muscle: 200 µg/kg

Fat or fat+skin: 300 µg/kg

Liver: 500 µg/kg

Kidney: 1500 µg/kg

Chicken:

Muscle: 400 µg/kg

Skin + fat: 250 µg/kg

Liver: 800 µg/kg

Kidney: 1000 µg/kg

**Manufacturer: Y.S.P. INDUSTRIES (M) SDN. BHD.**

Factory: Lot 3, 5 & 7, Jalan P/7, Section 13,  
Kawasan Perindustrian Bandar Baru Bangi,  
43000 Kajang, Selangor Darul Ehsan, Malaysia.  
Tel: 03-89251215 (4 Lines) Fax: 03-89251298

Office: No. 18, Jalan Wan Kadir, Taman Tun Dr. Ismail,  
60000 Kuala Lumpur, Malaysia.  
Tel: 03-77276390 (12 Lines)  
Ordering line: 1800 88 3027  
Product info: 1 800 88 3679

Date: 10 Jul 2009