

Dimension: L-160 mm & W-100 mm (Auto DS)

Kop[®] 2.5% Gel

Ketoprofen BP

1. NAME OF THE MEDICINAL PRODUCT

Kop[®] 2.5% Gel

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Ketoprofen 2.5% w/w

3. PHARMACEUTICAL FORM & DESCRIPTION

Gel; A clear transparent gel having odour of lavender oil, filled and sealed in a printed Laminated tube with polypropylene cap. Free from any undispersed particles.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

For Inflammatory non-articular musculoskeletal conditions and mild trauma due to sport injuries, sprains, tendinitis, musculotendinous contusion and oedema.

4.2 Posology and method of administration

Dosage: Adult : Apply twice daily to the skin in the painful or inflamed region for up to 7 days. Apply gently and massage well to ensure gel penetration. The total dose should not exceed 15g /day regardless of the number of areas affected. Do not dilute the gel. Children (under 15 years): Not recommended, as safety in children has not been established.

4.3 Contraindications

Patients with a known hypersensitivity to ketoprofen or any of the excipients, aspirin or other non-steroidal anti-inflammatory agents; patients suffering from or with a history of bronchial asthma or allergic disease. Ketoprofen gel should be avoided in patients with exudative dermatoses, eczema, sores and infected skin lesions or broken skin.

Ketoprofen gel should not be applied to mucous membranes, anal or genital areas, eyes or used with occlusive dressings. Ketoprofen is contraindicated during the last trimester of pregnancy.

4.4 Special warnings and precautions for use

Although systemic effects are minimal, the gel should be used with caution in patients with severe renal impairment. Should a skin rash occur after gel application, treatment must be stopped. Do not apply Ketoprofen gel beneath occlusive dressings. Areas of skin treated with Ketoprofen gel should not be exposed directly sunlight, or solarium ultraviolet light, either during treatment or for two weeks following treatment discontinuation. Keep the gel away from naked flames. Do not incinerate.

4.5 Interaction with other medicinal products and other forms of interaction

Interactions are unlikely as serum concentrations following topical administration are low.

Serious interactions have been recorded after the use of high dose methotrexate with non-steroidal anti-inflammatory agents, including ketoprofen, when administered by the systemic route.

4.6 Pregnancy and lactation

No embryopathic effects have been demonstrated in animals and there is epidemiological evidence of the safety of ketoprofen in human pregnancy.

Nevertheless, it is recommended that ketoprofen should be avoided during the first and second trimester of pregnancy.

During the third trimester of pregnancy, all prostaglandin synthetase inhibitors including ketoprofen may induce cardiopulmonary and renal toxicity in the fetus. At the end of the pregnancy, prolonged bleeding time in both mother and child may occur. Non-steroidal anti-inflammatory drugs may also delay labour. Therefore, ketoprofen is contraindicated during the last trimester of pregnancy.

Trace amounts of ketoprofen are excreted in breast milk, therefore Ketoprofen Gel should not be used during breast feeding.

4.7 Effects on ability to drive and use machines

No effect on drive or using machines.

4.8 Undesirable effects

Adverse reactions: Skin reactions, including photosensitivity reactions, pruritus, localised erythema. These are usually mild and resolve after cessation of the gel. Cases of more severe reactions such as bullous or phlyctenar eczema which may spread or become generalised have occurred rarely.

4.9 Overdose

Overdose is unlikely by topical administration.

If accidentally ingested, the gel may cause systemic adverse effects depending on the amount ingested. However, if they occur, treatment should be supportive and symptomatic.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Ketoprofen is a non-steroidal anti-inflammatory drug. It has antiinflammatory and analgesic actions.

5.2 Pharmacokinetic properties

Pharmacological data says that Plasma and tissue levels of ketoprofen have been measured in 24 patients undergoing knee surgery. After repeated percutaneous administration of Ketoprofen gel the plasma levels were about 60 fold less (9-39 ng/g) than those obtained after a single oral dose of ketoprofen (490-3300 ng/g). Tissue levels at the area of application were within the same concentration range for the gel as for the oral treatment, although the gel was associated with a considerably higher inter-individual variability.

The bioavailability of ketoprofen after topical administration has been estimated to be approximately 5% of the level obtained after an orally administered dose, based on urinary excretion data.

The protein binding in plasma is approximately 99%. Ketoprofen is excreted through the kidneys mainly as glucuronide conjugate.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Carbopol-940
- Triethanolamine
- Methyl Hydroxybenzoate
- Propyl Hydroxybenzoate
- Ethanol
- Menthol
- Lavendar oil
- Purified water

6.2 Incompatibilities

The gel should not be diluted.

6.3 Special precautions for storage

Store below 30°C. Protect from light.

Keep out of children's reach.

Jauhkan dari kanak-kanak.

6.4 Nature and contents of container

Laminated tube containing 20 gm gel.

Manufactured by :



SQUARE PHARMACEUTICALS PLC.

Square Road, Salgaria, Pabna 6600,
Bangladesh

© Registered Trade Mark

Date of revision : November 2023

2000026918