

ANUCARE® OINTMENT

Ingredient(s):

Each gram contains:

Zinc Oxide 107.5mg (10.75% w/w)

Peru balsam 18.8mg (1.88% w/w)

Benzyl Benzoate 12.5mg (1.25% w/w)

Preservative:

Methyl Paraben 1mg (0.1% w/w)

Propyl Paraben 1mg (0.1% w/w)

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetics):

Each gram of the Ointment contains Zinc Oxide 107.5mg, Peru Balsam 18.8mg and Benzyl Benzoate 12.5mg. Zinc Oxide is applied to the skin as a mild astringent, as a soothing and protective application in haemorrhoids. Peru Balsam has antiseptic action by virtue of its content of cinnamic and benzoic acids, it is used for the treatment of eczema, pruritus and symptomatic relief of haemorrhoids. Benzyl Benzoate has the antipruritic properties and the combination of Benzyl benzoate together with Zinc oxide and Peru Balsam is used in the treatment of uncomplicated haemorrhoids, simple anorectal inflammation and irritations caused by or associated with proctitis or cryptitis.

The active ingredients exert their therapeutic effect without being absorbed into the systemic circulation. Benzyl benzoate is not appreciably absorbed through the skin after topical application.

Indication(s):

For the relief of itching, burning and soreness of haemorrhoids and minor rectal inflammations and irritations.

Dosage and Administration:

Wash and dry affected area. Apply ointment twice daily ensuring area is completely covered. Attach the plastic applicator tip onto the tube. Insert tip into the rectum and gently squeeze the tube. After use wash nozzle in a hot solution of detergent and then thoroughly rinse in clear water.

Contraindication(s):

Benzyl benzoate is irritant to the eyes and mucous membranes and it may be irritant to the skin. Hypersensitivity reactions have been reported.

Hypersensitivity to any other components of the preparation.

Precaution(s) / Warning(s):

Consult a doctor if symptoms persist. Do not swallow.

Interaction with Other Medicaments

None.

Pregnancy and Lactation

There is no epidemiological evidence of adverse effect, either to the pregnant mother or fetus.

Until more information is available, this product should only be used during pregnancy or breastfeeding if the potential benefit of treatment outweighs potential risks.

Side Effect(s) / Adverse Reaction(s):

Sensitivity reactions are rare. Discontinue medication if suspected. Local irritation may be experienced.

Symptoms and Treatment for Overdosage, and Antidote(s):

None have been mentioned.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Keep in an airtight container. Store at temperature below 30°C. Protect from light and moisture.

Product Description & Packing(s):

Opaque brownish ointment.

Aluminium tube of 10g, 15g, 20g, 10g x 50, 15g x 40, 15g x 72, 20g x 40, 20g x 72 and 50g.



Manufacturer and Product Registration Holder:
Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)
Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.
Ordering Line: 1 800 88 3027
Product Info: 1 800 88 3679