

Name: Hirudoid®_insert (Malaysia)
Design artwork by BIO-LIFE VBC Team
Version : C1 - version 3
Date : 17/04/2015



Hirudoid® Cream

Mucopolysaccharide polysulphate 0.3g/100g

Product description

White, shining, homogeneous cream with a characteristic odor for cutaneous use only.

Pharmacodynamics

Chondroitin polysulfate (mucopolysaccharide polysulfate) has antithrombotic, fibrinolysis promoting and anti-inflammatory characteristics. Animal experiments and human-pharmacological studies have shown that, after topical application of the compound, there is a faster absorption of subcutaneous hemorrhages and fibrinates and a faster dissolution of superficial thrombi. An anti-inflammatory effect has been demonstrated in different experimental inflammation models.

Pharmacokinetics

A permeation of chondroitin polysulfate into superficial tissue was shown in animal experiments and in human skin with radioactively labelled active constituent, with chemical-analytical and histo-chemical methods. According to a specific concentration gradient, mucopolysaccharide spread into the tissue without any systemic effect. Blood coagulation could not be shown, even when the substance was applied for several days.

Indication

For the topical treatment of

- bruises with or without hematomas
- superficial thromboses and phlebitis
- inflammation of varicose veins.

Recommended dosage

Hirudoid® cream is to be applied to the affected area 2-3 times daily, if necessary, it may be used more frequently. Depending on the size of the area to be treated, a length of 3 to 5 cm of cream is usually sufficient.

Hirudoid® cream is massaged into the skin gently. In the case of painful inflammation, the cream is distributed carefully on the affected area. Hirudoid® cream is also suitable for cream dressings.

The primary therapeutic measure to be taken for superficial phlebitis of the lower extremities is compression treatment.

Avoid contact with the eyes, the mucous membranes, open wounds or injured skin.

The usual treatment period for blunt traumata is up to 10 days, one to two weeks for superficial phlebitis & two to six weeks for varicose veins.

Children: up to now, the use and safety of Hirudoid® cream in children and adolescents has not been reviewed systematically.

Contraindications
Hirudoid® must not be used if there is hypersensitivity to chondroitin polysulfate or to any of the excipients of the medicinal product.

Hirudoid® should not be applied to open wounds or injured skin.

Hirudoid® cream must not be used if there is a hypersensitivity to methyl-4-hydroxybenzoate or propyl-4-hydroxybenzoate.

Warnings and precautions
Cetostearyl alcohol and wool wax alcohol ointment may cause regional skin irritations (e.g. contact dermatitis).

Avoid applying to bleeding or infected areas.
Avoid contact with the eyes or mucous membranes.

Interactions with other medicaments

None known

Pregnancy and lactation

There are no reports on damaging effects on the unborn child or the infant when chondroitin polysulfate is used topically in pregnant or lactating women.

Adverse effects

The following frequencies are applied when evaluating side effects: very common ($\geq 1/10$), common ($\geq 1/100$), uncommon ($\geq 1/1000$, $< 1/100$), rare ($\geq 1/10,000$, $< 1/1000$), very rare/individual cases ($< 1/10,000$).

Skin and hypodermis disorders

In individual cases local hypersensitivity reactions, such as reddening of the skin, may occur, which however usually subside quickly after discontinuation of the product.

Immune system disorders

Methyl- and propyl-4-hydroxybenzoate may cause hypersensitivity reactions also of the late type.

Overdose and treatment

a) Symptoms of overdose

When applied as directed, an overdose of Hirudoid® is not known. If Hirudoid® is taken orally by accident, no signs of overdose caused by the active constituent are to be expected.

b) Therapeutic measures in case of overdose

Not applicable.

Storage condition

Keep in cool dry place below 30°C.

Packaging available:

5g, 14g and 40g per tube

Manufacturer

OLIC (Thailand) Limited
Bangpa-In Industrial Estate, Ayuthaya,
13160 Thailand.

For Medinova AG, Switzerland

Date of revision: 23 March 2015

medinova
SWITZERLAND

Pantone Red 032C