



For animals treatment only
Keep medicine out of reach of children/ Jauhi ubat daripada kanak-kanak

Planate™ Injection

Cloprostenol 87.5 µg/ml

Composition:

Each 2 ml dose contains 184 µg of Cloprostenol Sodium BP (Vet) equivalent to 175 µg of cloprostenol. The solution also contains 20.0 mg/ml Benzyl Alcohol as a preservative.

Product Description: Clear, colourless solution, practically free from particles.

Pharmacodynamics:

The Prostaglandin F_{2α} analogue cloprostenol has luteolytic activity. Following its administration plasma progesterone falls to baseline levels. As a consequence, parturition is initiated and proceeds normally. The effect of cloprostenol on the smooth muscular system is similar to that of Prostaglandin F_{2α} itself.

Pharmacokinetics:

Following injection, cloprostenol is rapidly absorbed and a peak plasma concentration of 1 ng/ml is reached within 8 min after injection. A very rapid elimination of cloprostenol then occurs until 1.5 hours, followed by a slower elimination phase leading to concentrations below quantifiable levels between 4 and 6 hours post-administration.

Indication:

A synthetic prostaglandin analogue for use in pigs. Planate is a luteolytic agent which induces farrowing in sows and gilts.

Dosage and Administration:

Single 2ml dose.

Administration: By deep intramuscular injection, observing normal aseptic precautions. It is recommended that a 1 ½ (4 cm) needle be used. Having calculated the average gestation length for each farm, sows and gilts may be injected two days before this date or on any date thereafter to suit the requirements of the particular management system. Trials have shown that normally 95% of animals will commence farrowing within 36 hours of treatment. The majority of animals can be expected to respond within the period 24 (N6) hours following injection, except in those cases where spontaneous farrowing is imminent.

Recommendation: Planate provides opportunity for more efficient and convenient management under a variety of systems. Allows batch management of sows and gilts to be efficiently achieved.

Minimises farrowing at weekends, public holidays and during the night. Facilitates supervision of farrowing. Facilitates interbreeding. Enables farrowing and labour schedules to be planned for convenience. Prevents sows and gilts going beyond term. Allows optimal use of farrowing quarters, equipment, etc.

Contraindication:

Do not use in pregnant animals, for which induction of abortion or parturition is not intended. Do not use in animals with spastic diseases of the respiratory or gastrointestinal tract.

Warnings/ Precautions:

Special warning for each target species

None.

Special precautions for use

- Special precautions for use in animals

Induction of farrowing too early in pregnancy can lead to non-viable piglets being born. An increase in the number of non-viable piglets may result if used more than two days prior to the average gestation length calculated from farm records.

- Special precautions to be taken by the person administering the product to animals

Direct contact with skin or mucous membranes of the user should be avoided.

Prostaglandins of the F_{2α} type may be absorbed through the skin and **may cause bronchospasm or miscarriage**. Care should be taken when handling the product to **AVOID SELF-INJECTION OR SKIN CONTACT**.

Pregnant women, women of child-bearing age, asthmatics and persons with other respiratory tract diseases should exercise caution when handling cloprostenol. Those persons should avoid contact or wear disposable plastic gloves during administration of the product.

Accidental spillage on the skin should be washed off immediately with soap and water. The possible incidence of bronchospasm with the product is unknown. Should shortness of breath result from accidental inhalation or injection, seek urgent medical advice and show the doctor this warning. Wash hands after use.

Pregnancy & Lactation: Do not use in pregnant animals, for which induction of abortion or parturition is not intended.

The safety of the veterinary medicinal product has not been established during lactation. There are no data suggesting negative effects of the treatment with cloprostenol on the course of lactation.

Symptoms & Treatment of Overdose: None known.

Adverse Effects: None known.

Interaction with Other Medicines: None known.

Incompatibilities: None known.

Withdrawal Period:

Animals must not be slaughtered for human consumption during treatment. Pigs may be slaughtered for human consumption only after 2 days from the last treatment.

Shelf life after first opening of container: 4 weeks

Storage Condition: Protect from light, store below 30°C

Package Quantities: 20 ml colourless tubular glass, type I. Each vial pack in a carton.

Manufactured by: Vet Pharma Friesoythe GmbH

A Division of Essex Pharma GmbH, Sedelsberger Strabe 2, 26169 Friesoythe, Germany.

Marketing Registration Holder: Intervet (M) Sdn. Bhd., B-22-1 & B-22-2, The Ascent Paradigm, No. 1, Jalan SS7 /26A, Kelana Jaya, 47301 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

Date of Revision of Package Insert: 25th January 2021