



Vaxcel®

SODIUM FUSIDATE 500mg INJECTION

COMPOSITION:

1 vial containing dry substance & 1 vial containing solvent. The dry substance for injection contains : Sodium Fusidate 500mg.

PRESENTATION:

A vial of white or almost white crystalline powder packed with clear solvent that provides a clear solution after reconstitution.

INDICATIONS:

Vaxcel Sodium Fusidate 500mg Injection is indicated for the treatment of infections caused by susceptible organisms, especially staphylococci, e.g. osteomyelitis, septicaemia, endocarditis, superinfected cystic fibrosis, pneumonia, skin and soft tissue infections, surgical and traumatic wound infections.

PHARMACODYNAMICS:

Pharmacology:

Sodium Fusidate is the sodium salt of Fusidic Acid, which is an antibiotic which mode of action is as inhibitor of the bacterial protein synthesis resulting in a bactericidal effect. Fusidic Acid has a powerful antibacterial activity against a number of gram-positive microorganisms.

Staphylococci, including the strains resistant to penicillin, methicillin or other antibiotics, are particularly susceptible to Fusidic Acid. Fusidic Acid shows no cross-resistance to any other antibiotic agent used in clinical practice. Fusidic Acid has an unusually good ability to penetrate tissue and it widely distributed in the organism. It is of great clinical importance that Fusidic Acid provides high concentrations not only in areas well supplied with blood, but also in relatively avascular tissue. Concentration exceeding the MIC for *Staphylococcus aureus* (0.03 - 0.16mcg/ml) have been found in pus, sputum, soft tissue, heart tissue, bone tissue, synovial fluid, sequestra, burn crusts, brain abscesses, and intraocularly.

PHARMACOKINETICS:

Fusidic Acid is metabolized in the liver and mainly excreted through the bile, little or none being excreted through the urine. Fusidic Acid is generally toxic and shows no cross-hypersensitivity to other antibiotics in clinical use which is why Fusidic Acid also can be used in many cases where other antibiotics are contraindicated, e.g. in patients allergic to penicillin or to other antibiotics.

In severe or deep-seated infections and when prolonged therapy is required, systemic Fusidic Acid should generally be given concurrently with other antistaphylococcal antibiotic therapy to minimize the risk of resistance development. Fusidic Acid may be combined with penicillinase-stable penicillins, cephalosporins, erythromycin, aminoglycosides, lincomycin, rifampicin or vancomycin, and thus an additive or synergistic effect is obtained.

DOSAGE AND ADMINISTRATION:

Vaxcel Sodium Fusidate 500mg Injection may be given as intravenous infusion whenever oral therapy is inappropriate, including cases where absorption from the gastrointestinal tract is unpredictable.

Adult: 500mg (1 vial) 3 times daily

Children: 20mg/kg/day divided in 3 equal doses.

According to the metabolism and excretion of Vaxcel Sodium Fusidate 500mg Injection, no dosage adjustment is needed in renal impairment or in patients undergoing hemodialysis as Vaxcel Sodium Fusidate 500mg Injection is not significantly dialysed.

ROUTE OF ADMINISTRATIONS:

Only for Intravenous infusion.

Dissolve 1 vial of Vaxcel Sodium Fusidate 500mg Injection in the 10ml sterile buffer solution (Vaxcel Phosphate Citrate Buffer) provided. Dilute to 250 - 500ml with Sodium Chloride Injection or 5% Dextrose Injection BP. The infusion solution should be used with 24 hours. Opalescence may be encountered with more acidic samples of dextrose and the solution should then be discarded. The infusion period should not be less than 2 - 4 hours. Vaxcel Sodium Fusidate 500mg Injection should be made into a wide-bore vein with good blood flow or through a central venous catheter to minimize the risk of venospasms and thrombophlebitis. For adults a total daily dose of 2g should not be exceeded.

The solution of Vaxcel Sodium Fusidate 500mg Injection in the buffer must never be injected undiluted. Due to local tissue injury Vaxcel Sodium Fusidate 500mg Injection must not be given intramuscularly or subcutaneously.

CONTRAINDICATIONS:

Hypersensitivity to Sodium Fusidate/ Fusidic Acid and other ingredients in the product.

SIDE EFFECTS:

Thrombophlebitis and venospasm may be seen when using Vaxcel Sodium Fusidate 500mg Injection for intravenous infusion. Reversible increase of transaminases is reported at daily dosage of 1.5-3g of Vaxcel Sodium Fusidate 500mg Injection. Reversible jaundice has been reported in some patients after administration of Vaxcel Sodium Fusidate 500mg Injection, most frequently in patients receiving intravenous Vaxcel Sodium Fusidate 500mg Injection in high dosage and especially in serious *Staph. aureus* bacteremia. In some cases, instituting oral therapy may be beneficial. If the jaundice persists, Vaxcel Sodium Fusidate 500mg Injection should be withdrawn, after which the serum bilirubin will return to normal. Allergic reactions are reported in very few cases.

OVERDOSAGE AND TREATMENTS:

There has been no experience of overdosage. Treatment should be restricted to symptomatic and supportive measures. Dialysis

is of no benefit, since the drug is not significantly dialysed.

WARNING AND PRECAUTION:

Due to the metabolism and excretion of Fusidic Acid, periodic liver function tests should be carried out in patients with liver dye-function, abnormalities in the biliary pathway, when Fusidic Acid is given in high doses for prolonged periods, or when it is given in combination with other antibiotics which have similar excretion pathways, e.g. lincomycin and rifampicin. In vitro, Fusidic Acid displaces bilirubin from its albumin binding site. The clinical significance of this finding is uncertain and kernicterus has not been observed in neonates receiving Fusidic Acid. However, this observation should be borne in mind when the drug is given to preterm, jaundiced acidotic or seriously ill neonates.

Pregnancy & Lactation:

Animal studies and many years of clinical experience suggest that Fusidic Acid is devoid of teratogenic effects. Fusidic Acid passes the placenta and should be avoided during third trimester due to the theoretical risk of kernicterus. Concentrations of Fusidic Acid found in breast milk are negligible and its use is not contraindicated in nursing mothers.

INTERACTION WITH OTHER MEDICAMENTS:

In few cases, Fusidic acid has been reported to potentiate the anticoagulant efficacy of coumarin derivatives.

The use of sodium fusidate systemic should be avoided in patients treated with CYP-3A4 biotransformed drugs. Coadministration of sodium fusidate systemic and HMG-CoA reductase inhibitors such as statins causes increased plasma concentrations of both agents resulting in an elevation of creatine kinase level (rhabdomyolysis), muscle weakness and pain. Coadministration of sodium fusidate systemic and HIV protease inhibitors such as Ritonavir and Saquinavir causes increased plasma concentrations of both agents which may result in hepatotoxicity. Coadministration of sodium fusidate systemic and Cyclosporin has been reported to cause increased plasma concentration of Cyclosporin.

Incompatibilities:

Vaxcel Sodium Fusidate 500mg Injection contains Sodium Fusidate which is incompatible with kanamycin, gentamicin, cancomycin, cephaloridine, carbenicillin, whole blood, amino-acid solutions and calcium-containing solutions. Precipitation of Fusidic Acid may occur in infusion solutions with pH values below 7.4.

STORAGE:

Store below 30°C. Protect from heat and direct sunlight. Reconstituted solution should be stored below 25°C and used within 24 hours.

**KEEP OUT OF REACH OF CHILDREN
JAUHI DARI KANAK-KANAK**

PACK QUANTITIES:

Each combination pack contains 2 vials. 1 vial of Vaxcel Sodium Fusidate 500mg Injection. 1 vial Vaxcel Phosphate Citrate Buffer as a solvent.

Further information can be obtained from pharmacist, physician or the manufacturer.

Product Registration Holder/ Manufacturer



Kotra Pharma (M) Sdn. Bhd.
1, Jalan TTC 12,
(90082-V)
Cheng Industrial Estate,
75250 Melaka, Malaysia.

MAL11110006AZ

REGN-090620