



PIASCLEDINE®

Capsule

code 128 : 8305918

Read this leaflet carefully before you start taking the medicine. Important information regarding your treatment is enclosed.

If you have further questions, or if in doubt, please ask your doctor or your pharmacist for additional information.

Keep this leaflet. You may need to read it again. Ask your pharmacist if you need more information or advice. You must see a doctor if your symptoms worsen or do not improve.

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet.

QUALITATIVE AND QUANTITATIVE COMPOSITION:

Total extract of avocado and soybean unsaponifiable fractions.....300.00mg

Including:

Unsaponifiable fraction of avocado oil100.00mg

Unsaponifiable fraction of soybean oil 200.00mg

Excipients: anhydrous colloidal silica, butylhydroxytoluene

Composition of the capsule shell: gelatin, polysorbate 80, titanium dioxide, erythrosine, yellow oxide of iron.

PHARMACEUTICAL FORM:

Capsule (orange and grey) marked "P300". Box of 30 or 60 capsules.

PHARMACOTHERAPEUTIC GROUP

OSTEOARTHRITIS:

Pharmacotherapeutic group: Other anti-inflammatory and anti-rheumatic agents, non-steroids.

ATC code: M01AX26

PIASCLEDINE® also belongs to the class of "SYSADOA" (symptomatic slow-acting drugs in osteoarthritis), which are notably characterized by their delayed action.

NAME AND ADDRESS OF MANUFACTURER:

Laboratoires Expanscience
Rue des quatre filles
28230 EPERNON-FRANCE

WHAT IS THE MEDICINE USED FOR?

PIASCLEDINE® is used for the symptomatic slow-acting treatment of knee osteoarthritis in adults.

The full efficacy of PIASCLEDINE® may be delayed by approximately 2 months and may persist for 1 to 2 months after treatment discontinuation. You must talk to a doctor if you do not feel better after using the medicine for approximately 2 months or if you feel worse.

CAUTION!

PLEASE CONSULT THE DOCTOR BEFORE USE IN ORDER TO EXCLUDE THE PRESENCE OF OTHER JOINT CONDITIONS OR DISEASES FOR WHICH AN ALTERNATIVE TREATMENT MAY BE REQUIRED.

WHEN THE MEDICINE IS NOT TO BE USED? (CONTRAINDICATIONS)

The medicine **MUST NOT BE TAKEN** if you are allergic to one of the active substances or any of the other ingredients of this medicine.

SPECIAL PRECAUTIONS FOR USE:

IF IN DOUBT, DO NOT HESITATE TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE.

KEEP THE MEDICINE OUT OF THE REACH OF CHILDREN.

If you suffer from or have history of liver or gallbladder disease, inform your doctor or pharmacist before taking PIASCLEDINE®. If you develop any signs or symptoms of liver or biliary problems, such as unexplained nausea, vomiting, abdominal pain, tiredness, loss of appetite, yellowing of the skin and eyes (jaundice) and dark urine, stop the treatment and contact immediately your doctor. Stop the treatment and contact immediately your doctor if you develop any signs and symptoms of allergic reactions such as itching, hives, skin rash/redness, swelling of the face, breathing difficulty.

Several cases of decreased platelet counts (thrombocytopenia) have been reported. Before taking PIASCLEDINE®, tell your doctor or pharmacist if you are treated with anticoagulants (see section "Interaction with other medicinal products and other forms of interaction").

Discontinue PIASCLEDINE® treatment and contact immediately your doctor if you develop any signs and symptoms of coagulation disorders such as red or purple spots on the skin, spontaneous bruising or bleeding into the mucous membranes.

PIASCLEDINE® is for adults only. Do not give this medicine to children and adolescents below the age of 18 years.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

TO AVOID ANY INTERACTION WITH OTHER MEDICINAL PRODUCTS, YOU HAVE TO TELL YOUR DOCTOR OR PHARMACIST IF YOU HAVE TAKING, HAVE RECENTLY TAKEN OR MIGHT TAKE ANY OTHER MEDICINES.

If you are receiving anticoagulants treatment, please consult your doctor before taking PIASCLEDINE® due to a potential risk of coagulation disorders (bleeding and/or bruising).

PREGNANCY - BREAST-FEEDING – AND FERTILITY

The use of PIASCLEDINE® should be avoided during pregnancy. PIASCLEDINE® should not be used during breast-feeding.

GENERALLY SPEAKING, DURING PREGNANCY OR BREAST-FEEDING, IT IS RECOMMENDED TO ASK YOUR DOCTOR OR PHARMACIST FOR ADVICE BEFORE TAKING ANY MEDICINE.

The use of PIASCLEDINE® is not recommended in women of childbearing potential with no contraception. There is not enough information on its effect on fertility.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

PIASCLEDINE® does not affect the ability to drive and use machines.

HOW TO USE THE MEDICINE

POSOLOGY

Always use this medicine as your doctor or pharmacist has told you. If you are unsure, you should check with your doctor or pharmacist.

The maximum recommended dose is 1 capsule daily, taken during a meal.

The effect of PIASCLEDINE® is delayed by approximately 2 months and it may persist for 1 to 2 months after treatment discontinuation.

Do not discontinue the treatment if you feel its efficacy does not appear quickly enough, before talking to a doctor or pharmacist. Do not exceed the recommended dose.

METHOD AND ROUTE OF ADMINISTRATION

Oral route. The capsule should be swallowed whole with a glass of water. Do not chew.

FREQUENCY AND MOMENT OF THE INTAKE

Everyday, during a meal.

For an adequate medicine use, you must comply with a regular medical follow-up.

IF YOU TAKE MORE PIASCLEDINE THAN YOU SHOULD

Consult immediately your doctor or pharmacist.

SIGNS AND SYMPTOMS OF OVERDOSE

Overdose of PIASCLEDINE® may induce or exacerbate gastrointestinal and/or hepatic disorders. If you have taken more than 1 capsule per day, consult immediately your doctor or pharmacist.

IF YOU FORGET TO TAKE PIASCLEDINE

Do not take a double dose to make up for a forgotten dose. Take the next planned dose of the medicine.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

POSSIBLE UNWANTED SIDE EFFECTS

(Undesirable effects)

LIKE ALL ACTIVE MEDICINES, PIASCLEDINE® CAN CAUSE SIDE EFFECTS IN SOME PERSONS. The most frequent side effects with PIASCLEDINE® are diarrhoea, abdominal pain, heartburn, nausea and upper abdominal discomfort.

Possible serious side effects:

Tell your doctor immediately and stop taking this medicine if you have any of the following serious side effects which may require urgent medical treatment:

Sign and symptoms that may be a sign of severe allergic reactions (uncommon frequency): Itching, hives, skin rash/redness, swelling of the face, breathing difficulty.

Intense skin eruption (rare frequency)

Hepatic disorders (uncommon frequency): Liver damage or bile ducts obstruction both sometimes associated with yellow skin, as well as increased level of liver enzymes in blood (such as transaminases, alkaline phosphatase, bilirubin and gamma-glutamyl transpeptidase) may occur with uncommon frequency (see section "special precautions for use").

The following other side effects have also been reported:

- Common (may affect up to 1 in 10 people): Diarrhea
- Uncommon (may affect up to 1 in 100 people): Upper abdominal discomfort, heartburn, nausea, bad taste in the mouth (that could be avoided by taking the capsule during a meal), abdominal pain, discoloured faeces, headache, dark urine, weakness, tiredness, breast pain, breast swelling, gynaecological bleeding
- Rare (may affect up to 1 in 1,000 people): Gastrointestinal disorders, such as inflammation of bowels, vomiting, dry mouth, eructation, eczema, kidney stones, coagulation disorders such as decreased platelets count (see section "special precautions for use"), blood pressure increase

IF YOU GET ANY UNWANTED SIDE EFFECTS, TALK TO YOUR DOCTOR OR PHARMACIST. THIS INCLUDES ANY POSSIBLE SIDE EFFECTS NOT LISTED IN THIS LEAFLET.

STORAGE AND DISPOSAL:

DO NOT TAKE THIS DRUG AFTER THE EXPIRY DATE WHICH IS CLEARLY STATED ON THE PACK (THE EXPIRY DATE REFERS TO THE LAST DAY OF THE MONTH).

Keep this medicine out of the sight and reach of children.

Store below 30°C.

Do not use this medicine if you notice any visible signs of deterioration.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use.

FOR ANY INFORMATION ABOUT THIS MEDICINE, PLEASE CONTACT THE MARKETING AUTHORISATION HOLDER:

Hyphens Pharma Sdn. Bhd.
C-L2-01, Block C, Axis Business Park,
No. 10, Jalan Bersatu 13/4,
46200 Petaling Jaya, Selangor, Malaysia

DATE OF THE REVISION OF THE TEXT: 06/2025

1. NAME OF THE MEDICINAL PRODUCT ***** OK DEF *****

PIASCLEDINE® capsule.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

2.1 General Description

PIASCLEDINE® capsule is made of soybean and avocado

vegetal extracts.

2.2 QUALITATIVE AND QUANTITATIVE COMPOSITION

Each capsule contains 300mg of avocado and soybean oil unsaponifiables, corresponding to:

- Avocado oil unsaponifiable : 100.00 mg,
 - Soybean oil unsaponifiable : 200.00 mg.
- For a full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Hard capsule, size No1, orange body and opaque greyish cap, containing a soft brown paste and marked "P 300".

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

PIASCLEDINE® capsule is indicated for the symptomatic slow-acting treatment of knee osteoarthritis in adults.

4.2 Posology and method of administration

Posology

Adult Patients:

The recommended dose is one capsule once daily. This daily dose should not be exceeded. Increased dosing in clinical trials did not result in enhanced clinical benefit. The duration of the treatment will be adapted according to the clinical results, knowing that the effect is delayed by approximately 2 months. The effect may persist for 1 to 2 months after discontinuation of the treatment.

Paediatric Population:

In the absence of efficacy or safety data in children and adolescents below 18 years, the use of PIASCLEDINE® is not recommended in this population.

Elderly Patients:

No dosage adjustment is necessary for elderly. In the clinical studies conducted with PIASCLEDINE® the population mean age was above 60 years, with a significant proportion of patients over 65 years.

Renal Impairment:

No data are available in patients with pre-existing renal impairment. Therefore, no recommendation can be made.

Hepatic Impairment:

No data are available in patients with pre-existing severe hepatic impairment. Therefore, no recommendation can be made.

In case of worsening of hepatic or biliary impairment after PIASCLEDINE® initiation, PIASCLEDINE® treatment should be discontinued (see section 4.4).

Method of administration

Oral use. The capsule is to be swallowed whole, without chewing, with a glass of water, during a meal.

4.3 Contraindications

Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

During the post marketing period, adverse reactions such as hepatic cytolysis, cholestasis, jaundice, and transaminases increase were reported (see section 4.8). Therefore, this information must be taken into consideration before initiating treatment with PIASCLEDINE® in patients with a medical history or suffering from hepatic or biliary disorders or presenting any medical condition that may increase risk of cholelithiasis or hepatic injury. PIASCLEDINE® should be discontinued in case of biological or clinical symptoms of hepatic or biliary injury.

Hypersensitivity reactions as well as cutaneous rash, urticaria, dermatitis were reported during the clinical development of the product (see section 4.8). Patients should be advised of signs and/or symptoms suggestive of an allergic reaction and should stop the treatment and consult their physician at the first occurrence of a cutaneous reaction or signs of hypersensitivity.

Few cases of thrombocytopenia have been reported with PIASCLEDINE® including one during the clinical development. PIASCLEDINE® should be discontinued in case of symptoms indicative of coagulation disorders (e.g. petechia, purpura).

4.5 Interaction with other medicinal products and other forms of interaction

PIASCLEDINE® should be taken with caution in patients treated concomitantly with anticoagulants due to potential risk of coagulation disorders (see sections 4.4 and 4.8.)

4.6 Fertility, pregnancy, and lactation

Pregnancy

There are no clinical data on the use of PIASCLEDINE® in pregnant women.

The use of PIASCLEDINE® should be avoided during pregnancy.

Breast-feeding

There are no clinical data on the use of PIASCLEDINE® in breast feeding women and it is unknown whether PIASCLEDINE® or its metabolites are excreted in human milk. Therefore, PIASCLEDINE® should not be used during breast-feeding.

Fertility

Signs of reproductive toxicity have been shown in animals receiving high doses of PIASCLEDINE® (see section 5.3.).

There is no clinical data to evaluate PIASCLEDINE® effect on human fertility.

The use of PIASCLEDINE® is not recommended in women of childbearing potential with no contraception.

4.7 Effects on ability to drive and use machines

PIASCLEDINE® has no or negligible influence on the ability to drive and use machines.

4.8 Undesirable effects

1. Summary of the safety profile

The most frequently reported adverse drug reactions during PIASCLEDINE® therapy are related to gastro-intestinal disorders occurring in approximately 3% of patients. Diarrhoea, abdominal pain, dyspepsia, nausea, abdominal distension may occur between 0.4% to 1.1% of patients. These adverse reactions were mild or moderate in intensity and were dose dependent.

2. Tabulated list of adverse reactions

Adverse reactions reported during clinical trials and controlled clinical trials versus placebo / active comparator including 1,310 patients treated with PIASCLEDINE® (1,174 of these patients were treated with PIASCLEDINE® 300 mg once daily) and/or adverse reactions reported during the post marketing use are listed in the table below. Tabulated summary of adverse drug reactions by body system, frequency [Very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥1/10,000 to <1/1,000); very rare (<1/10,000)]. Within each frequency category, serious reactions are presented first:

System Organ Class	Frequency	Adverse Reactions
Blood and lymphatic system disorders	Rare	Thrombocytopenia
Immune system disorders	Uncommon	Hypersensitivity (*)
Nervous system disorders	Uncommon	Headache
Gastrointestinal disorders	Common	Diarrhoea
	Uncommon	Dyspepsia; Dysgeusia (S); Nausea; Abdominal pain; Discoloured faeces (**)
	Rare	Enteritis; Vomiting; Eructation; Dry mouth
Hepatobiliary disorders (#) (**)	Uncommon	Hepatic cytolysis; Cholestasis; Jaundice; Increased transaminases, alkaline phosphatase, bilirubin and gamma glutamyl transpeptidase
Skin and subcutaneous tissue disorders	Rare	Toxic skin eruption (*); Eczema (*)
	Uncommon	Chromaturia (**)
Renal and urinary disorders	Rare	Nephrolithiasis
	Uncommon	Breast pain; Breast swelling; Metrorrhagia
Reproductive system and breast disorders (#)	Uncommon	
General disorders and administration site conditions	Uncommon	Asthenic conditions
Investigations	Rare	Increased blood pressure (***)

(S) The capsule should be taken during a meal to avoid taste disorders. See section 4.2.

(#) These adverse reactions have been reported during the post marketing period. As such reactions were not reported during clinical trials including 1,310 patients

composed of PIASCLEDINE®, their frequency is estimated as uncommon.

Description of selected adverse reactions

(*) Intake of PIASCLEDINE® may in some cases lead to occurrence of a skin allergic reaction or hypersensitivity such as systemic allergic reactions, i.e. acute onset of an illness with involvement of the skin (pruritus, urticaria, erythema, rash), mucosal tissue, or both, cough, persistent gastrointestinal symptoms, or reduced blood pressure and/or associated symptoms. See section 4.4.

(**) Signs of hepatic disorders such as abdominal pain associated with nausea, faeces discoloration, chromaturia and/or icterus may occur with PIASCLEDINE® treatment. See section 4.4.

(***) Signs and symptoms suggestive of blood pressure increased (i.e. headache, hearing disorders, flushing) may occur with PIASCLEDINE® treatment.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via <https://www.npra.gov.my/index.php/en/health-professionals/reporting-adr>.

4.9 Overdose

Overdose (intake of more than 1 capsule a day) may induce or exacerbate gastro intestinal disorders and/ or hepatic disorders. Such disorders, if they occur, may be symptomatically treated and PIASCLEDINE® treatment should be reconsidered.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group:

Other anti-inflammatory and anti-rheumatic agents, nonsteroids. ATC code: M01AX26.

Pharmacodynamic effects

With trophic target

Clinical efficacy and safety

The effects of Piascledine 300 mg in osteoarthritis have been evaluated in clinical studies and have shown positive results on the onset of symptoms. The results of five randomized controlled studies (three placebo-controlled studies and two active comparator studies) are presented below.

In a multicentre, randomized, double-blind, placebo-controlled clinical study on 86 patients with osteoarthritis of the knee, the consumption of NSAIDs was significantly reduced by 70% compared to 33% for the placebo (p <0.01) after 3 months. Pain at rest and algofunctional disorders were also significantly improved with Piascledine 300 mg, compared to placebo.

In another multicentre, randomized, double-blind, placebo-controlled clinical study in a subgroup of 101 patients with knee osteoarthritis, the primary endpoint was the resumption of NSAIDs after discontinuation. The number of patients resuming NSAID consumption over 3 months was 54.2% with Piascledine 300 mg, compared to 73% with the placebo (p = 0.05).

A significant decrease in algofunctional disorders was observed, while there was no significant decrease in pain at rest.

In a 6-month, multicentre, randomized, double-blind, placebo-controlled clinical study Piascledine 300 mg did not significantly improve the primary endpoint (Lequesne index, p = 0.07) or pain at rest in the subgroup of patients with knee osteoarthritis (N = 55) possibly due to the small number of patients, but significantly decreased the consumption of NSAIDs.

Two non-inferiority studies compared Piascledine 300 mg with chondroitin sulfate 400 mg three times daily for 6 months and diacerin 50 mg twice daily for 3 months in 184 and 64 patients respectively. These trials showed that Piascledine 300 mg was non-inferior to both comparators on the total WOMAC score and on the consumption of NSAIDs, which were the primary endpoint of each study respectively. No difference was observed between these treatments and Piascledine 300 mg regarding the secondary endpoints (Lequesne index and pain at rest).

In addition, Piascledine 300 mg is slow acting, delayed by about 2 months and its effect may persist after treatment discontinuation.

5.2 Pharmacokinetic properties

PIASCLEDINE® is an extract composed of a multitude of entities from soya and avocado origin for which no validated analytical methods are available. Therefore, no pharmacokinetic investigations were conducted.

5.3 Preclinical safety data

The general toxicological properties of PIASCLEDINE® have been studied in various animal species, including rodents, rabbits, and dogs, in acute, subacute, and chronic dosing conditions.

Acute and sub-chronic toxicity studies have shown a low toxicity of PIASCLEDINE®, with no LD50 calculated due to the absence of lethality for doses up to 8000 mg/kg. Some liver and thyroid abnormalities were shown in rats treated with the highest dose (750 mg/kg).

In chronic toxicity studies (6 months) in rats and dogs, the general tolerance of PIASCLEDINE® was satisfactory; however, liver and thyroid were also confirmed as target organs. Moderate liver and thyroid laboratory and histopathological abnormalities were shown in the medium and high doses groups, more pronounced in dogs than in rats. In both species, the low PIASCLEDINE® dose (30 – 50 mg/kg, depending on the species) was associated with minimal toxicity.

PIASCLEDINE® is not a metabolic inducer in rat. A series of repro-toxicity studies were performed with PIASCLEDINE®.

- Fertility:
 - o No effect on male rat fertility
 - o In female rats, at 750 mg/kg/day, some adverse findings on implantation and/or very early survival of embryos were reported.
- Embryotoxicity: in rats and rabbits slight skeletal modifications were reported at high doses (750 mg/kg and 500 mg/kg, respectively). Skeletal observations in rabbits at 50 and 150 mg/kg were not dose-related and difficult to interpret in relation to the treatment. A confirmatory study performed in rabbit in order to clarify those abnormalities did not reveal any relevant skeletal modification. At 500 mg/kg, PIASCLEDINE® treatment augmented the number of post-implantation loss. Altogether, the NOAEL values for embryotoxicity were established at 200 mg/kg (rat) and 150 mg/kg (rabbit).
- Peri and post-natal toxicity: No effect on F0 and F1 generations.

Mechanistic (other toxicology studies) studies confirmed that in rats the liver was a target organ for PIASCLEDINE®, at least partly through its avocado unsaponifiable fraction. The NOAEL was 10 mg/kg/day for the avocado unsaponifiable fraction. Studies conducted with PIASCLEDINE® did not show any mutagenic or genotoxic effects.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Capsule content:

Anhydrous colloidal silica (Aerosil 200), Butylhydroxytoluene (E321)

Capsule shell:

Polysorbate 80, Gelatine, Titanium dioxide (E171), Erythrosin (E127), Yellow oxide of iron (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years

6.4 Special precautions for storage

Store below 30°C.

6.5 Nature and contents of container

Heat-formed PVC/PVDC blister and aluminium foil.

Cardboard box containing 1, 2, or 4 blisters of 15 capsules.

Not all pack sizes are available.

6.6 Special precautions for disposal

Do not throw away any medicines via wastewater or household waste as these measures will help to protect the environment. Recommend returning to specified pharmacies for disposal.

7. MANUFACTURER

Laboratoires Expanscience

Rue des quatre filles 28230 EPERNON-France

8. MARKETING AUTHORISATION HOLDER

Hyphens Pharma Sdn. Bhd.

C-L2-01, Block C, Axis Business Park, No. 10, Jalan Bersatu 13/4, 46200 Petaling Jaya, Selangor, Malaysia.