



LION BALL MEDICATED OIL

Composition: Methyl Salicylate 25% w/v, Menthol 16% w/v, Camphor 5% w/v, Eucalyptus Oil 20 % w/v, white oil up to 100 % w/v

Description: A transparent colour oily liquid with menthol smell, packed in a glass bottle.

Indication: For temporary relief of minor aches and pains of muscle and joint associated with arthritis, strains and sprains.

Direction of Use: Apply the oil on affected part with rubbing, 3 – 4 times daily.

Route of Administration: Topical

Pharmacodynamics

Camphor is a stimulant, rubefacient, antipruritic and feebly antiseptic. When in combination with other ingredient like menthol, methyl salicylate, it becomes ideal for neuralgia and other painful areas.

Menthol dilates the blood vessels causing a sensation of coldness, followed by an analgesic effect. Menthol also acts as a penetration enhancer, increasing the penetration of drugs when applied on the skin, to give a faster onset of action.

Methyl Salicylate is a salicylic acid derivative. Salicylates inhibit cyclooxygenase, thereby reducing the formation of prostaglandins and cause platelet dysfunction. Methyl salicylate is used topically as a counter irritant. Upon application, it is absorbed through the skin and is applied for the relief of the pain in rheumatic conditions and painful muscle or joints.

Pharmacokinetics

Camphor is readily absorbed from all administration sites. Camphor is well absorbed after dermal exposure. It is hydroxylated in the liver to yield hydroxycamphor metabolites which are then conjugated with glucuronic acid and excreted in the urine. Camphor crosses the placenta.

After absorption, *menthol* is excreted in the urine and bile as a glucuronide.

Methyl Salicylate may be absorbed through intact skin. The absorption of topical salicylates is proportional to the surface area involved, duration of exposure, concentration and skin integrity. Absorption characteristics of salicylates vary with the dose, formulation and route of administration. Percutaneous absorption is enhanced by exercise, heat, occlusion or disruption of the integrity of the skin or application to large areas of skin. Both the rate and extent of absorption increases after repeated application, increasing the bioavailability. Methyl salicylate is extensively metabolised to salicylic acid in the dermal and subcutaneous tissues after topical application. At therapeutic levels, the half-life of salicylates is 2– 4 hours. As salicylate level increase into the toxic range, the half- life can be greater than 18 hours.

Contraindications:

Do not use in patients with aspirin or salicylate idiosyncrasy.

Patient with known hypersensitivity to any of the ingredients in the product.

Warnings and Precautions:

This product is contraindicated in children below 2 years of age. Caution must be exercised when older children are treated. It is dangerous to place any camphor containing product into the nostril of children. A small amount applied this way may cause immediate collapse.

Topical analgesic preparations containing methyl salicylate should be used with caution in patients at an increased risk of developing salicylate adverse effects.

Initially apply a small amount to a limited skin area and observe whether there is discomfort or itch before applying it to a larger skin area.

Discontinue use if irritation develops. Avoid contact with the eyes and mucous membranes. Do not apply to open wounds or damaged skin. Avoid excessive use during pregnancy and lactation. Do not apply over large areas of the body or bandage or use with heating pads. If symptoms persist, consult your doctor.

For external use only. Keep out of reach of children.

Effects on Ability to Drive and Use Machines: Not known

Interactions with Other Medicaments: Warfarin

Statement on Usage During Pregnancy and Lactation: Safety for use in pregnancy has not been established; therefore the potential benefit of the product should be weighed against the possible risks to the mother and child.

Adverse Effects/ Undesirable Effects:

If severe irritation such as redness and burning sensation occurs, discontinue use immediately and consult a physician.

Symptoms and Treatment of Overdose:

Symptoms of methyl salicylate overdose

Salicylate intoxication can occur after ingestion or topical application of methyl salicylate. Mild chronic salicylate intoxication or salicylism, usually occurs only after repeated use of large doses. Salicylism can also occur following excessive topical application of salicylates. Symptoms include dizziness, tinnitus, deafness, sweating, nausea and vomiting, headache and confusion and may be controlled by reducing the dosage.

Symptoms of more severe intoxication or of acute poisoning following overdose include hyperventilation, fever, restlessness, ketosis and respiratory alkalosis and metabolic acidosis. Depression of the CNS may lead to coma; cardiovascular collapse and respiratory failure may also occur.

Symptoms of camphor/ menthol overdose

Ingestion of significant quantities is reported to cause symptom: severe abdominal pain, nausea, vomiting, vertigo, ataxia, drowsiness and coma; instant collapse in infants after the local application of menthol or camphor to their nostrils.

Seizures may be the first clinical sign of severe toxicity of camphor; however, seizures are usually self- limited. Severe toxicity of camphor can result in delirium, visual hallucinations, cerebral edema and status epilepticus. Systemic toxicity may include hypotension, tachycardia, respiratory failure and death.

Treatment

The stomach should be emptied by gastric lavage or administration of oral activated charcoal. Fluid and electrolyte management is the mainstay of treatment with the immediate aim of correction of acidosis, hyperpyrexia, hypokalemia and dehydration if present.

Any convulsion must be controlled first through supportive care including anticonvulsant therapy.

Storage Conditions: Store at room temperature below 30°C.

Shelf Life: Please refer to the packaging. Do not use beyond the expiry date.

Dosage Form: Liniment

Packing: 3 ml, 5 ml, 10 ml, 45 ml, 56 ml, 90 ml

The information contained in this leaflet is limited. For further information, please consult your doctor or pharmacist.

Product Registration Holder:

GAB Sdn Bhd (Co. No. 863197A)
No. 24, Jalan BPU 4, Bandar Puchong Utama,
47100 Puchong, Selangor.

Manufacturer:

Wellmetech Pharmaceutical Industries Sdn Bhd
Lot 8058, Samarahan Industrial Park,
94300 Kota Samarahan, Sarawak, Malaysia.

Date of Revision: Oct 2024



MINYAK ANGIN BOLASINGA

Komposisi: Metil Salisilat 25% w/v, Mentol 16% w/v, *Camphor* 5% w/v, Minyak Kayu Putih 20 % w/v, *white oil* tambah ke 100 % w/v
Deskripsi: Cecair berminyak warna jernih dengan bau mentol, dibungkus dalam botol kaca.

Indikasi: Untuk kelegaan sementara sakit otot dan sendi yang ringan akibat artritis, terseliuh dan ketegangan.

Cara Penggunaan: Sapukan pada bahagian yang terabit sambil menggosok 3 - 4 kali sehari.

Laluan Pemberian Ubat: Topikal

Farmakodinamik: Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik: Sila rujuk kepada penerangan Bahasa Inggeris.

Kontraindikasi: Jangan gunakan pada pesakit dengan idiosyncrasi aspirin atau salicylate. Pesakit yang diketahui hipersensitif kepada mana-mana ramuan dalam produk.

Amaran dan Perhatian:

Ubat ini kontraindikasi pada kanak-kanak di bawah umur 2 tahun. Perhatian perlu diberikan kepada kanak-kanak berumur 2 tahun ke atas apabila menggunakannya. Sapu ubat yang mengandungi camphor pada hidung kanak-kanak adalah berbahaya kerana boleh menyebabkan pengsan.

Persediaan analgesik topikal yang mengandungi metil salisilat harus digunakan dengan berhati-hati pada pesakit pada peningkatan risiko mengembangkan efek salisilat.

Pada mulanya sapu kuantiti yang kecil pada kawasan kulit yang terhad dan memerhati sama ada ketidakselesaan atau gatal berlaku sebelum sapu pada kawasan kulit yang lebih besar. Jika iritasi kulit berlaku, sila berhenti menggunakannya. Elakkan bersentuhan dengan mata dan membran mukus.

Jangan sapu pada luka terbuka atau kulit yang rosak. Elakkan penggunaan yang berlebihan semasa kehamilan dan penyusuan. Jangan gunakan di bahagian besar badan atau pembalut atau gunakan dengan pad pemanasan. Sekiranya gejala berterusan, merujuk kepada doktor anda.

Untuk kegunaan luar sahaja. Jauhi daripada capaian kanak-kanak.

Kesan ke atas Keupayaan Memandu dan Penggunaan Mesin: Tidak diketahui

Interaksi dengan ubat lain: Warfarin

Kehamilan dan Penyusuan: Keselamatan untuk kegunaan semasa kehamilan belum dimantapkan; Oleh itu potensi manfaat produk terhadap kemungkinan risiko kepada ibu dan anak harus ditimbang.

Kesan Sampingan: Jika kerengsaan seperti kemerahan dan rasa terbakar berlaku, hentikan penggunaan serta merta dan rujuk kepada doktor.

Simptom dan Rawatan untuk Kelebihan Dos: Sila rujuk kepada penerangan Bahasa Inggeris.

Kedudukan Penyimpanan: Simpan pada suhu bilik kurang daripada 30°C.

Jangka Hayat: Sila rujuk kepada bungkusan. Jangan guna selepas tamat tarikh luput

Bungkusan: 3 ml, 5 ml, 10 ml, 45 ml, 56 ml, 90 ml

Maklumat terkandung dalam risalah ini adalah terhad. Untuk maklumat terperinci, sila rujuk kepada ahli farmasi atau doktor anda.



狮球风油

成份: 冬青油 25% w/v · 薄荷 16% w/v · 樟脑 5% w/v · 桉树油 20 % w/v · 白油添加至 100 % w/v

描述: 带有薄荷气味的透明油状液体 · 装在玻璃瓶中。

主治: 用于暂时缓解与关节炎、扭伤和拉伤引起的肌肉和关节的轻微疼痛。

用法: 搽在患处并揉搓 · 每日 3 至 4 次。

给药途径: 外用

药效学: 请参阅英文版本。

药代动力学: 请参阅英文版本。

禁忌: 禁用于对阿司匹林或水杨酸盐有特异性反应的患者。对产品中任何成分已知过敏的患者禁用。

警告和注意事项:

本产品禁用于 2 岁以下儿童。用于较年长的儿童时需多加谨慎。将含有樟脑的产品涂抹于儿童的鼻腔内是很危险的，此法少量应用可能会导致立即晕厥。

对于服用华法林的患者，因肌肉或关节疼痛而过度涂抹在皮肤上可能会增加出血的机会。

最初应将少量药物涂于一小块皮肤区域，观察是否有不适或瘙痒，然后再涂抹于更大面积的皮肤。如果出现刺激反应，应立即停止使用。避免接触眼睛和黏膜。不要涂抹于开放性伤口或受损的皮肤。

孕和哺乳期间应避免过度使用。不要在身体大面积区域使用或包扎后使用本产品，也不要与加热垫一起使用。如果症状持续，请咨询医生。

仅供外用。请放置于儿童无法触及的地方

对驾驶和操作机器能力的影响: 未知

与其他药物相互作用: 华法林

妊娠和哺乳期间使用说明: 本产品在妊娠期间使用的安全性尚未确立；因此，应权衡其潜在的益处与对母婴可能照成的风险。

副作用: 如果出现严重的刺激反应，如发红和灼热感，请立即停止使用并咨询医生。

过量症状和治疗: 请参阅英文版本。

储藏: 置于在 30°C 以下的室温环境中。

有效期: 请参见包装说明。勿使用超过有效期的产品。

剂型: 搽剂

包装: 3 毫升、5 毫升、10 毫升、45 毫升、56 毫升、90 毫升

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。