

Prazitel™ Plus+Tablets For Dogs and Puppies

Statement of the active substance(s) and other ingredient(s): 1 tablet contains 50 mg praziquantel, 144 mg pyrantel embonate and 150 mg febantel.

Product description: Yellow coloured, uncoated, circular, flat bevelled tablets with cross breakline on one side and plain on the other.

Pharmacodynamics: This product contains anthelmintics active against gastrointestinal roundworms and tapeworms. The product contains three active substances, as follows:

- 1. Febantel, a probenzimidazole
- 2. Pyrantel embonate (pamoate), a tetrahydropyrimidine derivative
- 3. Praziquantel, a partially hydrogenated pyrazinoisoquinoline derivative

In this fixed combination, pyrantel and febantel act against all relevant nematodes (ascarids, hookworms, and whipworms) in dogs. In particular, the activity spectrum covers *Toxocara canis*, *Toxascaris leonina*, *Uncinaria stenocephala*, *Ancylostoma caninum* and *Trichuris vulpis*. This combination shows synergistic activity in the case of hookworms and febantel is effective against *T. vulpis*.

The spectrum of activity of praziquantel covers all important cestode species in dogs, in particular *Taenia spp.*, *Dipylidium caninum*, *Echinococcus granulosus* and *Echinococcus multilocularis*. Praziquantel acts against all adult and immature forms of these parasites.

Pyrantel acts as a cholinergic agonist. Its mode of action is to stimulate nicotinic cholinergic receptors of the parasite, induce spastic paralysis of the nematodes and thereby allow removal from the gastrointestinal system by peristalsis.

Within the mammalian system, febantel undergoes ring closure, forming fenbendazole and oxfendazole. It is these chemical entities which exert the anthelmintic effect by inhibition of tubulin polymerisation. Formation of microtubules is thereby prevented, resulting in disruption of structures vital to the normal functioning of the helminth. Glucose uptake in particular is affected, leading to a depletion in cell ATP. The parasite dies upon exhaustion of its energy reserves, which occurs 2 - 3 days later.

Praziquantel is very rapidly absorbed through the parasite's surface and distributed throughout the parasite. Both *in vitro* and *in vivo* studies have shown that praziquantel causes severe damage to the parasite integument, resulting in the contraction and paralysis of the parasites. There is an almost instantaneous titanic contraction of the parasite musculature and a rapid vacuolization of the syncytial tegument. This rapid contraction has been explained by changes in divalent cation fluxes, especially calcium.

Pharmacokinetics: Perorally administered praziquantel is absorbed almost completely from the intestinal tract. After absorption, the drug is distributed to all organs. Praziquantel is metabolized into inactive forms in the liver and secreted in bile. It is excreted within 24 hours to more than 95% of the administered dosage. Only traces of non-metabolised praziquantel are excreted.

Following administration of the product to dogs, peak plasma concentrations of praziquantel were achieved by approximately 2.5 hours.

The pamoate salt of pyrantel has low aqueous solubility, an attribute that reduces absorption from the gut and allows the drug to reach and be effective against parasites in the large intestine. Following absorption, pyrantel pamoate is quickly and almost completely metabolized into inactive metabolites that are excreted rapidly in the urine.

Febantel is absorbed relatively rapidly and metabolized to a number of metabolites including fenbendazole and oxfendazole, which have anthelmintic activity.

Following administration of the product to dogs, peak plasma concentrations of fenbendazole and oxfendazole were achieved by approximately 7-9 hours.

Indications: Tablets for the control of tapeworms, roundworms, hookworms and whipworms in dogs and puppies.

Worms controlled include:

Ascarids: *Toxocara canis* and *Toxascaris leonina* (adult and late immature forms).

Hookworms: *Uncinaria stenocephala* and *Ancylostoma caninum* (adults).

Whipworms: *Trichuris vulpis* (adults).

Tapeworms: *Echinococcus* species, *Taenia* species and *Dipylidium caninum* (adults and immature forms).

Contraindications/Interaction: Do not use simultaneously with piperazine compounds.

Adverse reactions: None Known. If you notice any serious effects or other effects not mentioned in this leaflet, please inform your veterinary surgeon.

Target species: Dogs and Puppies

Dosage for each species, routes and method of administration

For oral administration only.

The recommended dose rates are: 15mg/kg bodyweight febantel, 5 mg/kg pyrantel (equivalent to 14.4 mg/kg pyrantel embonate) and 5 mg/kg praziquantel. This is equivalent to 1 Prazitel Plus tablet per 10 kg (22 lbs) bodyweight.

The tablets can be given directly to the dog or disguised in food. No starvation is needed before or after treatment.

If there is a risk for re-infestation, the advice of a veterinarian should be sought regarding the need for and the frequency of repeat administration.

Special precautions for use in animals: Any part used tablet should be discarded.

Consult a veterinary surgeon before treating pregnant animals for roundworms.

Do not exceed the stated dose when treating pregnant bitches.

Special precautions for the disposal of unused product or waste material: Any unused product or waste material should be disposed of in accordance with national requirements. Medicines should not be disposed of via wastewater or household waste.

Ask your veterinary surgeon how to dispose of medicines no longer required. These measures should help to protect the environment.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Fleas serve as intermediate hosts for one common type of tapeworm – *Dipylidium caninum*. Tapeworm infestation is certain to reoccur unless control of intermediate hosts such as fleas, mice, etc. is undertaken.

In the interests of good hygiene, persons administering the tablets directly to the dog, or by adding them to the dog's food, should wash their hands afterwards.

Statement on usage during pregnancy and lactation: The product may be used during lactation. For the control of *Toxocara*, nursing bitches should be dosed 2 weeks after giving birth and every two weeks until weaning.

Overdose and Treatment: The combination of Praziquantel, Pyrantel Embonate and Febantel is well tolerated in dogs. In safety studies, doses of 5 times the recommended dose or greater gave rise to occasional vomiting.

Special storage precautions: Do not use after expiry date.

Do not store above 25°C. Protect from light.

Withdrawal period: N/A

Other Information: 104 tablets.

Manufactured by :

chanelle
Chanelle Pharmaceuticals Manufacturing Ltd.,
Loughrea, Co. Galway, Ireland.

Registration holder and repacker:

Rhone Ma
RHONE MA MALAYSIA SDN BHD (200001023270)
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

5LPRAZI3-104 04/XXX
Date of revision: February 2021

W10cm H14cm

LABEL CODE:	VERSION	COLORS
5LPRAZI3-104		BLACK
PRODUCT: PRAZITEL PLUS TABLET	Version:	
PACK SIZE: 104 TABLETS	MMM/YY:	
COUNTRY: MY		
DIMENSIONS: cm = 10 X 14		
		! Please use the official PANTONE® (P) matching system for an accurate color representation.

Marketing approval:	Regulatory approval:
COMPLETE NAME:	COMPLETE NAME:
DATE + SIGNATURE:	DATE + SIGNATURE: