

VETOQUINOL	Nom du produit : Tolfédine 6MG MY
Créé par : P.JARDEL	Référence article : 1018 A L
le :	Dimensions : 160 x 120 mm
Visa :	1 couleur
	

Diarrhoea and vomiting may occur in rare cases during the treatment. Moreover, a temporary increase of thirst and/or diuresis may occur. In most of the cases, these signs cease spontaneously after the treatment. In case of overdose, administer symptomatic treatment

USER WARNINGS

In case of accidental contact with eyes, wash with plenty of water.

**TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION
KEEP OUT OF REACH OF CHILDREN / JAUHI DARIPADA
KANAK-KANAK
STORE IN A DRY PLACE - DO NOT STORE ABOVE 25° C
KEEP BLISTER STRIPS IN OUTER CARTON**

Manufactured by:

VETOQUINOL

Magny Vernois

70200 Lure (France)

To be supplied only on veterinary prescription

Registration Holder:

Gladron Chemicals Sdn. Bhd.

No.7, Jalan TP7,

UEP Industrial Park,

40400 Shah Alam,

Selangor Darul Ehsan, Malaysia

DATE OF REVISION: october 2018.

Dispose of used packaging in the household refuse.

Unused product should be returned to the veterinary surgeon.

For animal treatment only.

1018 A



Veterinary Pharmaceuticals
70200 LURE (France)



TOLFEDINE® TABLETS 6 MG

Non steroidal anti-inflammatory - analgesic - antipyretic

Each tablet contains: 6 mg tolfenamic acid active ingredient

White convex tablets with average mass of 0.08 g

Tablets are packed in PVC/Aluminium blisters.

Box of 16 tablets (2 blisters of 8 tablets)

Box of 20 tablets (2 blisters of 10 tablets)

INDICATIONS

Cats: In febrile syndrome (abscess, fever of unknown origin)

Dogs: Treatment for alleviation of acute episodes of inflammation and pain in chronic locomotor disease

PHARMACODYNAMIC PROPERTIES

Tolfenamic acid (N-(2-methyl-3-chlorophenyl) anthranilic acid) is a non-steroidal anti-inflammatory drug belonging to the fenamate group. Tolfenamic acid possesses anti-inflammatory, analgesic and antipyretic properties.

The anti-inflammatory activity of tolfenamic acid is especially due to an inhibition of cyclooxygenase leading to a reduction of prostaglandin synthesis and thromboxanes, major inflammatory mediators.

PHARMACOKINETIC PROPERTIES

Absorption:

In the dog, tolfenamic acid is rapidly absorbed. After a single oral administration of 4 mg / kg tolfenamic acid, the mean maximal plasma concentration (C_{max}) of about 4 µg / mL is reached in about 1 hour. When the same intake of tolfenamic acid is taken with food, C_{max} is 2 ± 3 µg / mL. These variations are due to a strong enterohepatic recycling of the product.

In the cat, the absorption is very fast. After a single oral administration of 4 mg / kg tolfenamic acid, the mean maximal plasma concentration (C_{max}) of about 5.6 µg / mL is reached in about 1 hour.

Distribution, metabolism, excretion

Tolfenamic acid is distributed in all organs with a strong concentration in plasma, digestive tract, liver, lungs and kidneys. The concentration in the brain however is low.

Tolfenamic acid and its metabolites cross little the placenta barrier.

Tolfenamic acid is excreted mainly unchanged and in a little part under the form of non active metabolites.

In dogs with renal insufficiency, the elimination of tolfenamic acid is unchanged.

DOSAGE AND ADMINISTRATION

Tablets must be administered by oral route with food.

The recommended dosage is 4 mg tolfenamic acid / kg bodyweight by oral administration with food once daily for 3 days according to the following table:

	Bodyweight (kg)												
	1 - 2	2.5 - 3.5	3.5 - 4.5	4 - 6	6-8	8-10	10-12.5	12.5 -15	7- 10	10- 20	20- 25	23- 35	35- 40
Number of 6mg Tablets	1	2	3										
Number of 20mg Tablets				1	1½	2	2½	3					
Number of 60mg Tablets									½	1	1½	2	2½

In dogs, subject to clinical response, the administration may be repeated every 7 days, ie 3 days of medication followed by 4 days without medication.

CONTRA-INDICATIONS

Do not administer to animals suffering from cardiac, hepatic or renal disease, where there is a possibility of gastro-intestinal ulceration

or bleeding or where there is evidence of a blood dyscrasia or hypersensitivity to the product.

WARNINGS AND PRECAUTIONS

NSAIDS can cause inhibition of phagocytosis and hence in the treatment of inflammatory conditions associated with bacterial infections appropriate concurrent antimicrobial therapy should be instigated.

Do not exceed the prescribed dosage or duration of treatment.

In dogs, long term treatment of over 3 months duration should be under regular veterinary supervision. In particular, dogs with hepatic insufficiency should be closely monitored.

Use in animals less than 6 weeks of age, or in aged animals, may involve additional risk. If such a use cannot be avoided, animals may require a reduced dosage and careful clinical management.

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

Concurrent administration of potential nephrotoxic drugs should be avoided.

It is preferable that the product is not administered to animals undergoing general anaesthesia until fully recovered.

Animals suffering from a chronic renal insufficiency and requiring an anti-inflammatory treatment may be treated with tolfenamic acid without requiring an adjustment of the dosage. However, the use of this product is contra-indicated in acute cases of renal insufficiency.

Interactions

Do not administer NSAIDs concurrently or within 24 hours of each other. Some NSAIDs may be highly bound to plasma proteins and compete with other highly bound drugs, which can lead to toxic effects. Do not administer in conjunction with glucocorticosteroids.

Use in pregnancy

Although studies in laboratory animals did not show any effect on reproduction, it is not advisable to administer the product during gestation.

Undesirable effects and overdose

In case of undesirable effects (anorexia, vomiting, diarrhoea, presence of blood in faeces) occurring during the treatment, your veterinarian should be contacted for advice.