

Packing:
Aluminum aerosol container (Available pack size i.e. 36g/75g/112g)

Symptoms and Treatment of Overdose
Topical administration of Relispray does not cause any toxic effects if used according to the recommended dosage.

However, salicylate intoxication can occur after ingestion or topical application of methyl salicylate (an active ingredient in Relispray) under certain conditions. Ingestion of methyl salicylate poses the threat of severe, rapid-onset salicylate poisoning because of its liquid concentrated form and lipid solubility. It is readily absorbed from the gastrointestinal tract and most is rapidly hydrolysed to free salicylate. The symptoms, which may appear within 2 hours of ingestion, are similar to those of salicylate poisoning in general, although methyl salicylate is expected to be more toxic because of its lipid solubility. Like other salicylates, methyl salicylate may be absorbed through intact skin. Percutaneous absorption is enhanced by exercise, heat, occlusion, or disruption of the integrity of the skin. The amount absorbed will also be increased by application to large areas of skin. Results from a study show that both the rate and extent of absorption are increased after repeated application.

Effects on ability to drive and use machines
No adverse effect known.

Storage Condition
Store below 30°C. Do not freeze. Avoid storage in direct sunlight or heat. Keep away from eyes.

Keep out of reach of children.

Shelf Life:
36 months

Revision date:
February 2025

Manufactured by
MIDASCARE PHARMACEUTICALS PVT. LTD.
B-16, MIDC, Wakuj, Chhatrapati
Sambhajinagar - 431136, India.

SS14L0694

RELISPRAY®
PAIN RELIEF SPRAY

Name and Strength of Active Substance	
Cinnamon oil (Dakchini ka tel)	0.5
Camphor (Karpura)	10.0
Clove oil (Lavang ka tel)	1.0
Eucalyptus oil (Nilgiri ka tel)	6.0
Menthol (Pudina ka phool)	4.0
Methyl Salicylate (Wintergreen tel)	20.0
Turpentine Oil	10.0
EXCIPIENTS	
Isopropyl Alcohol	7.5
Propylene Glycol	1.0

Product Description
A clear colorless liquid spray with characteristic odour, filled in an Aluminum aerosol container equipped with valve.

Indication
Used topically as a rubefacient for the symptomatic relief of pain in musculoskeletal, joint and soft-tissue disorders.

Recommended Dosage
Spray a thin layer of the product over affected area 3-4 times daily. Relispray® is used topically on the affected area by spraying.

Direction for use
Shake well before each use. Spray the contents from 5cm distance onto the affected area. Cover sprayed area with a cloth, to improve effectiveness.

Pharmacodynamic
Methyl salicylate is irritant to the skin and is used topically in rubefacient preparations in musculoskeletal, joint and soft tissue

disorders and for minor peripheral vascular disorders. It is readily absorbed through intact skin and can produce effects typical of systemic salicylates.

Applied externally, camphor acts as a rubefacient and mild analgesic. Applied externally clove oil is irritant but can produce local anesthesia. Menthol when applied to the skin, menthol dilates the blood vessels, causing a sensation of coldness followed by an analgesic effect. It relieves itching and is used in creams, lotion or ointments in pruritus and urticaria. Eucalyptus oil has slightly astringent effect. Cinnamon oil has slightly astringent effect.

The ingredient as mentioned above has also a counter irritant effect which is also responsible in relieving the pain when applied locally.

Pharmacokinetic
None Stated.

Route of Administration
Topical

Contraindications
The preparation must not be applied to the non intact skin or open wound.

Special Warning and Precautions Warning
This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding. This product is contraindicated in infants below 2 years of age. Caution must be exercised when older children are treated.

Precaution
It is dangerous to place any camphor containing product into nostril of children. A small amount applied this way may cause immediate collapse. Precaution must be taken while spraying Relispray that the spray does not come in contact with very sensitive skin such as eyelids, lips etc. Avoid contact with eyes or open wounds or non intact skin. If reddening of the skin occurs or irritation persists which is non tolerable then discontinue the use. This product contains methyl salicylate and when applied or rub on to the skin, can be absorbed through the skin into the blood. For patients taking warfarin, excessive application on to the skin for muscle or joint pains may increase the chances of bleeding. Liquid petroleum gas (propellant) is highly flammable.

Interaction with other medicaments
Please refer to precautions for possible interaction with warfarin.

Pregnancy and Neonates
Adverse reaction are not known or reported, if used during pregnancy, hence its safe uses during pregnancy could not be documented. Being a strong counter irritant preparation must not be used in the case of neonates.

Adverse effects
Not known when applied locally. Wintergreen oil is highly toxic. There are basically three adverse skin reaction that one may encounter when using essential oils.

1. Irritation of the skin.
2. Contacts allergy or sensitization.
3. Photo toxicity.

Folding: 2 vertical folds
Dimension: 105 x 90 mm (WxH)
Folded size: 35 x 90 mm