

Vétoquinol	Nom du produit : Notice Tolfédine 4% Malaysia	
Créé par : A.Milleret	Référence article : XXXXXX 0611 B	
le :	Dimensions : 160x120 mm	
Visa :	1 couleur	rev B
281 U		

TO BE SUPPLIED ONLY ON VETERINARY PRESCRIPTION
KEEP OUT OF REACH OF CHILDREN / JAUHI DARIPADA KANAK-KANAK
PROTECT FROM LIGHT - DO NOT STORE ABOVE 25° C

FURTHER INFORMATION

Follow up treatment may be continued by the use of Toldefine Tablets.

REGISTRATION NUMBER: MALXXXX

Manufactured by:
VETOQUINOL SA
Veterinary Pharmaceuticals
MAGNY-VERNOIS
70200 LURE (France)

Registration Holder
Gladron Chemicals Sdn. Bhd.
No.7, Jalan TP7,
UEP Industrial Park,
40400 Shah Alam,
Selangor Darul Ehsan.

NATURE AND CONTENTS OF CONTAINER

Cardboard box with 1 amber glass vial of 10 ml
Cardboard box with 1 amber glass vial of 30 ml
Not all pack sizes may be marketed.

For animal treatment only

XXXXXX 0611 B



Vétoquinol



TOLFEDINE® 4% W/V SOLUTION FOR INJECTION
FOR USE IN DOGS AND CATS

A clear, colourless to slightly yellow solution for injection.

Non steroidal anti-inflammatory - analgesic - antipyretic

Each ml contains tolfenamic acid 40mg as active ingredient with benzyl alcohol 10.4mg and sodium formaldehyde sulphonylate dihydrate 5mg as preservative.

Uses

Cats: Adjuvant treatment of upper respiratory disease in association with anti-microbial therapy

Dogs: Inflammatory and painful post-operative syndromes

Pharmacodynamic properties

Tolfenamic acid (N-52-methyl-3-chloropentyl) anthranilic acid) is belonging to the fenamate group. Substances of this group exert antiinflammatory, analgesic and antipyretic activities. Fenamates are classified as non steroidal antiinflammatory drug (NSAID), the antiinflammatory activity of tolfenamic acid is mainly due to an inhibition of cyclooxygenase and thus a reduction of the synthesis of prostaglandins, important inflammatory mediator.

Pharmacokinetic particulars

The pharmacokinetics of tolfenamic acid have been investigated in laboratory animals, in man and in the target species, dogs and cats.

Absorption

In the dogs, tolfenamic acid is readily absorbed either by oral or by injectable administration. By oral route, Cmax of about 4 µg / ml is attained after about 1 hour at the single dose of 4 mg / kg. When administering at the same dose during the meal tolfenamic acid Cmax is of 2 ± 3 µg /ml. These variation can be accounted for a greater enterohepatic recycling when administered with food. By injectable route, the maximum plasma concentration of about 4 µg / ml (s.c.) and about 3 µg / ml (i.m.) are obtained 2 hours after administration of 4 mg / kg i.m. as well as s.c.

In the cats, the absorption is quite rapid. By oral route, after a dose of 4 mg/kg a mean maximum concentration of about 5.6 µg / ml is attained . The mean Tmax is observed after 1 hour. By injectable route, a peak of 3.9 µg / ml is obtained within 1 hour after injection of 4 mg / kg.

Distribution

In the dogs and in the cats, tolfenamic acid is bound at over 99% to the plasma proteins.

Biotransformation

The metabolic fate of tolfenamic acid has been studied in rat, rabbit, dog and man. The extent of metabolism is depending on the species concerned.In the rat, the man and the rabbit, the main metabolites are two hydroxy-metabolites, on the contrary, in the dog, there is no formation of hydroxy-metabolites, only tolfenamic acid and its conjugate with glucuronic acid are found in urine.

Elimination

The hydroxylated metabolites and their conjugates are mainly excreted by the kidneys. The unchanged tolfenamic acid and its glucuronides are predominantly excreted into the bile. Moreover, tolfenamic acid undergoes an intensive enterohepatic recycling.

DOSAGE AND ADMINISTRATION

The recommended dose is 4 mg/kg. This corresponds to a single injection of 1ml/10kg, repeated once after 24 hours if required (1ml/10kg) or a single injection of 1ml/10kg, the treatment being continued by the oral route, using tablets.

In dogs administer by intramuscular or subcutaneous injection. For the treatment of post-operative pain, this is best given pre-operatively, either at the time of premedication or induction of anaesthesia.

In cats administration is by the subcutaneous route only.

CONTRA-INDICATIONS

Tolfenamic acid is contra-indicated in case of cardiac disease
Do not use in animals with impaired hepatic.function or acute renal insufficiency
Tolfenamic acid is contra-indicated in case of ulceration or digestive bleeding, in case of blood dyscrasia or hypersensitivity to tolfenamic acid.
Do not inject intramuscularly in cats.

SPECIAL WARNINGS FOR EACH TARGET SPECIES

NSAIDS can cause inhibition of phagocytosis and hence in the treatment of inflammatory conditions associated with bacterial infections appropriate concurrent antimicrobial therapy should be instigated

The use of insulin-type needle/syringe is advisable particularly in low-weight animals to ensure an accurate dose.

SPECIAL PRECAUTIONS FOR USE IN ANIMALS

Use in animals less than 6 weeks of age, or in aged animals, may involve additional risk. If such a use cannot be avoided animals may require a reduced dosage and careful clinical management is essential. Reduced metabolism and excretion in these animals should be considered.

Avoid use in any dehydrated, hypovolaemic or hypotensive animal, as there is a potential risk of increased renal toxicity.

Concurrent administration of potential nephrotoxic drugs should be avoided.

It is preferable that the product is not administered to animals undergoing general anaesthesia until fully recovered.

Do not exceed the prescribed dosage or duration of treatment. The scale of pain relief after pre-operative administration may be influenced by the severity and duration of the operation.

Animals suffering from a chronic renal insufficiency and requiring an anti-inflammatory treatment may be treated with tolfenamic acid without requiring an adjustment of the dosage. However, the use of this product is contra-indicated in acute cases of renal insufficiency.

In case of undesirable effects (anorexia, vomiting, diarrhoea, presence of blood in faeces) occurring during the treatment, your veterinarian should be contacted for advice and the possibility of stopping treatment should be considered.

USER WARNINGS

Take care to avoid accidental self-injection. In case of eye or skin contact, wash immediately with water.

ADVERSE REACTIONS (FREQUENCY AND SERIOUSNESS)

A temporary increase in thirst and/or diuresis may occur. In most of the cases, these signs cease spontaneously after treatment. Diarrhoea and vomiting may occur during treatment, where either persists treatment should be discontinued.
Local injection site reactions have been reported following administration of this product.

USE DURING PREGNANCY OR LACTATION

Do not treat pregnant animals.

OVERDOSE

In case of overdose, administer a symptomatic treatment.

INTERACTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

Do not administer other NSAIDs concurrently or within 24 hours of each other. Some NSAIDs may be highly bound to plasma proteins and compete with other highly bound drugs, which can lead to toxic effects.
Following withdrawal of the first dose, use the product within 28 days. Discard unused material.
Dispose of any unused product and empty containers in accordance with guidance from your local waste regulation authority.

INCOMPATIBILITIES

None known.