

Dicetel® 50 mg

Film-coated tablets

Dicetel® 100 mg

Film-coated tablets

50 or 100 mg pinaverium bromide



Read this entire leaflet carefully before you start taking this medicine.

Keep this leaflet. You may need to read it again. If you have questions not answered by this pamphlet, please ask your doctor or pharmacist. This medicine has been prescribed to you personally and you should not pass it on to others. It may harm them, even if their symptoms are the same as yours.

Dicetel film-coated tablets are orange coloured film-coated tablets for oral administration, engraved with "50" or "100" on one side (depending on the tablet strength) and "S" on the other side. Each tablet contains 50 mg or 100 mg of pinaverium bromide which is equivalent to 43.25mg or 86.5mg Pinaverium respectively.

Excipients (non-medicinal ingredients):

Core: microcrystalline cellulose, pre-gelatinised starch, lactose monohydrate, anhydrous colloidal silica, talc, magnesium stearate.

Coating: basic butylated metacrylate copolymer, sodium laurylsulfate, stearic acid, talc, microcrystalline cellulose, aluminium lake "sunset yellow" (E110), titanium dioxide (E171), hydroxypropyl methylcellulose.

Indications

- Treatment and relief of symptoms associated with irritable bowel syndrome (IBS): abdominal pain, bowel disturbances and intestinal discomfort.
- Treatment of symptoms related to functional disorders of biliary tract.

Dosage and administration

Always take Dicetel exactly as your doctor has prescribed. If you have any questions, contact your doctor or pharmacist.

If you forget to take your tablet(s), do not take a double dose to compensate for it. If you require further information, please ask your doctor or pharmacist for advice.

Adults:

• Dicetel 50 mg, fi lm-coated tablets:

The usual adult dosage is one tablet three times a day (total daily dose of 150mg). The dosage may be increased up to a maximum of 2 tablets taken three times a day. (maximum total daily dose of 300 mg)

• Dicetel 100 mg, fi lm-coated tablets:

The usual adult dose age is 1 tablet two times a day (total daily dose of 200 mg). The dosage may be increased up to a maximum of 1 tablet taken three times a day (maximum total daily dose of 300 mg).

Method of administration

Swallow the tablets without chewing or sucking them. Take the tablets with a glass of water in the middle of a meal.

The tablet should not be swallowed when in the lying position or just before bedtime.

Paediatric population:

The experience in paediatric patient (<18 years of age) is limited.

Contraindications

Do not take Dicetel if you are hypersensitive to the active substance or to any of the excipients.

Warnings and special precautions for use

Because of a risk of oesophageal lesions, instructions and the methods of administration should be carefully adhered to. Patients with pre-existing oesophageal lesion and/or hiatus hernia should pay special attention to correct application of Dicetel.

Dicetel should not be administered for the relief of motility dysfunction due to underlying organic disease.

There is only limited experience with the use of Dicetel in children and adolescents. Therefore, this medicine should not be given to them but to adults only.

This medicinal product contains lactose. Patient with rare hereditary problems of galactose intolerance, the Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Dicetel contains Sunset yellow, an azo colouring agent that may cause allergic reactions.

Interactions with other medications

Clinical trials have demonstrated the absence of any interaction between pinaverium bromide and digitalis drugs (heart medications), oral antidiabetics, insulin, oral anticoagulants and heparin.

Co-administration of an anticholinergic drug may enhance spasmolysis.

No interference with laboratory tests for drug level detection was observed.

Pregnancy and lactation

Ask your doctor or pharmacist for advice before taking any medicine during pregnancy.

Use in pregnancy

No clinical data on exposed pregnancies are available. Animal studies are insufficient with respect to effects on pregnancy, embryonal/foetal development, parturition (giving birth) and postnatal (following birth) development. The potential risk for humans is unknown. The safety of Dicetel during pregnancy has not been established. Consequently, in the pregnant patient, this drug should only be administered if, in the judgment of the physician, its use is essential to the welfare of the patient. Furthermore, the presence of bromine should be taken into account. Theoretically, administration of

pinaverium bromide at the end of pregnancy can affect the newborn neurologically (hypotony, sedation).

No such case however has been reported with Dicetel.

Dicetel should not be used during pregnancy unless your doctor decides it is clearly necessary.

Use during lactation

There is insufficient information on the excretion of pinaverium bromide in human and animal breast milk. Physico-chemical and available pharmacodynamic/toxicological data on Dicetel point to excretions in breast milk and a risk to the suckling infant cannot be excluded. Dicetel should not be used during breast-feeding.

Effects on ability to drive and use machines

No studies on the effects on the ability to drive and use machines have been performed.

Important information about the ingredients

If you have been told by your doctor that you have an intolerance to some sugars, especially lactose, glucose or galactose, contact your doctor before taking this medicinal product.

This product contains sunset yellow (E110) as an excipient, which may cause allergic reactions.

Undesirable effects

Like all medicines, Dicetel may cause side effects, although not everybody experiences them. If you notice any side effects not mentioned in this leaflet, or if any of the side effects gets serious, please inform your doctor or pharmacist.

The following adverse events have been reported spontaneously during post-marketing use. A precise frequency can not be estimated from available data (not known).

Gastrointestinal disorders

Gastro-intestinal disturbances have been observed, e.g. abdominal pain, diarrhea, nausea, vomiting, and dysphagia (difficulty swallowing).

Oesophageal lesion may occur when not applied as advised.

Skin and subcutaneous tissue disorders

Cutaneous (skin) effects have been observed, e.g. rash, pruritus (itchiness), urticaria (hives), and erythema (reddening of the skin).

Immune system disorders

Hypersensitivity reactions may occur.

Overdose

Currently there is no specific information on overdose related adverse reactions. No specific antidote is known; symptomatic treatment is recommended.

Pharmacodynamics

The following is a detailed description of how the active ingredients of Dicetel work. For further explanations please consult your doctor.

Pharmacotherapeutic group: Other drugs for functional bowel disorders Pinaverium bromide is an antispasmodic selectively acting on the gastro-intestinal tract. It is a calcium antagonist which inhibits the influx of calcium into intestinal smooth muscle cells. In

animals, pinaverium directly or indirectly reduces the effects of the stimulation of the sensitive afferences. It is free from anticholinergic effects. It is also devoid of effects on the cardiovascular system.

Pharmacokinetics

The following is a detailed description of how the active ingredients of Dicetel are metabolized by the body. For further explanations please consult your doctor.

After oral administration pinaverium bromide is rapidly absorbed with peak plasma concentrations occurring within one hour. The drug is extensively metabolised and eliminated via the liver. The elimination half-life is 1.5 hours.

Absolute bioavailability for the oral formulation is very low (< 1%). Major route of excretion is via the faeces. Plasma protein binding of pinaverium bromide is high (95-96%).

Preclinical Safety Data

The oral LD50 in animals ranges from 153.6 to 2153 mg/kg body weight. Toxicity of pinaverium bromide after oral administration was low. Signs of toxicity were mostly limited to general signs of toxicity, gastrointestinal symptoms and CNS symptoms. Pinaverium bromide did not display genotoxic or carcinogenic properties and had no teratogenic potential. At higher dose levels pinaverium bromide decreased pregnancy performance, but no relevant effect on pre- or post-natal development. Placental transfer of pinaverium bromide was not studied.

Based on toxicity studies, the lowest NOAEL was 300, 100 and 25 mg/kg for mice, rats and dogs, representing 50, 20 and 5 multiples of the human dose, respectively.

Incompatibilities

Not applicable.

Shelf life and storage conditions

- This product can be stored for up to 36 months.
- Do not use the medicine after the expiry date stated on the carton.
- Do not store above 30 °C.
- Store in the original package and keep the blister in the outer carton in order to protect from light.
- Keep this medicine out of the reach and sight of children.

Pack sizes

Dicetel 50mg: 15, 20, 25, 30, 40, 50, 60, 100 or 120 tablets per pack (not all pack sizes may be marketed). The blisters strips are made from PVC/aluminium.

Dicetel 100mg: 10, 15, 20, 25, 30, 50 or 100 tablets per pack (not all pack sizes may be marketed). The blisters strips are made from PVC/aluminium.

Further information

Any unused product or waste material should be disposed of in accordance with local requirements.

The information in this leaflet is limited. For further information, please contact your doctor or pharmacist.

Date of information

February 2012

Manufactured by:

Mylan Laboratories SAS
01400 Châtillon-sur-Chalaronne - FRANCE

Product Holder:

Abbott Laboratories (M) Sdn. Bhd.,
No. 22, Jalan Pemaju U1/15, Seksyen U1,
HICOM-Glenmarie Industrial Park,
40150 Shah Alam,
Selangor Darul Ehsan
Malaysia