

FRONT LABEL:

NUOQIU 0.5% W/W POWDER



White powder
0.5% (w/w) of Diclazuril

25.00 kg (in weight)

Manufactured By:

Shandong Luxi Animal Medicine Share Co., Ltd.
West of Qihe County, Shandong, P. R. China.

Batch Number:

Manufacturing Date :

Expiration Date :

Country's Registration No.: MALXXX

Registration Holder:

Gladron Chemicals Sdn. Bhd.
No.7, Jalan TP7,
UEP Industrial Park,
40400 Shah Alam,
Selangor Darul Ehsan,
Malaysia.

Keep out of reach of children/ Jauhi daripada kanak-kanak

Keep in cool, dry place at 25 °C and humidity of 60% RH.

Controlled Medicine/ Ubat Terkawal

For animal use only

Warnings / Precautions:

- As with all medicated feedstuffs, operators are advised to use protective clothing, gloves, dust mask and goggles when mixing or handling Nuoqiu 0.5% Powder.
- Do not use diclazuril medicated feed for the treatment of outbreaks of coccidiosis.
- If unexpected deaths occur, obtain an accurate diagnosis and follow veterinarian's or poultry pathologist's recommendations.
- Do not use with pellet binding agents.

Dosage and Administration:

> To be mixed in the feed.

> Sheep:

- 1mg diclazuril per kg bodyweight or 200mg of product per 2.5kg bodyweight at about 4-6 weeks of age at the time that coccidiosis can normally be expected on the farm.

- Under conditions of high infection pressure, a second treatment may be indicated about 3 weeks after the first dosing.

> Calves:

- 1mg diclazuril per kg bodyweight or 200mg of product per 2.5kg bodyweight, administered as a single dose, 14 days after moving into a potentially high risk environment.

> Chicken (Broilers):

- Use 0.0001% diclazuril medicated feed as the sole ration, from one day of age up until slaughter.

> Turkey (growing)

- Use 0.0001% diclazuril medicated feed as the sole ration, from one day of age to a maximum of 14 weeks of age.

- To obtain the recommended level of 0.0001% of diclazuril in the complete feed, the product should be mixed thoroughly with the feed at the rate of 200 grams per metric tonne. It is recommended that a premix be made from the product before final mixing. It is suggested to premix 200 grams of the product with 20-50 kg of a feed ingredient before incorporating in the feed.

BACK LABEL:

Withdrawal Period:

- Chicken (broiler): 5 days
- Turkey (growing): 0 days
- Sheep & Calves (meat & offal): 0 days

Pharmacodynamics:

- Diclazuril is an anticoccidial of the benzeneacetonitrile group and has anticoccidial activity against *Eimeria* species. Depending on the coccidian species, diclazuril has a coccidiocidal effect on the asexual or sexual stages of the development cycle of the parasite. Diclazuril treatment will only have limited effect on the intestinal lesions caused by parasitic stages older than 16 days. Treatment with diclazuril causes interruption of the coccidial cycle and of excretion of oocysts for approximately 2 weeks. This allows the animal to bridge the period of decrease of maternal immunity (observe data approximately 4 weeks of age).

Pharmacokinetics:

- The absorption of diclazuril in lambs is poor after administration of the oral suspension. Maximum concentrations in plasma are reached about 24 hours after dosing. The absorption decreases with the animals' age. The elimination half life is about 30 hours. In-vitro studies on sheep hepatocytes demonstrated that metabolic transformation of diclazuril is limited. This was equally observed in other animal species. Excretion occurs almost completely via the faeces.

Indication:

- In sheep: Prevention of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.
- In calves: Prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.
- In poultry: Chicken (broiler): Prevention of coccidiosis caused *Eimeria acervulina*, *E. maxima*, *E. tenella*, *E. brunetti*, *E. mitis*, *E. necatrix*.
- Turkey (growing): Prevention of coccidiosis caused by *Eimeria gallopavonis*, *E. adenoids*, *E. meleagrimitis*.

Contraindications:

- None known

Interactions with Other Medicaments:

- Not to mixed or used simultaneously with any other anticoccidial.

Pregnancy and Lactation

- Do not use in laying birds

Adverse Effects/ Undesirable Effects

- None known

Overdose and Treatment

- Diclazuril was administered to chickens at the maximum authorized level in feedingstuffs (1 mg/kg), and was well tolerated with no toxicity observed. Identical conclusions were drawn from experiments performed on female and male breeding broilers, fed with a diet at 5 mg/kg of diclazuril for 28 and 42 days respectively. Finally, at 25 times (25 mg/kg) the maximum authorized concentration of 1 mg/kg, chickens for fattening exposed from birth to day 37 showed no signs of intolerance. This was concluded to indicate a very large safety margin for diclazuril in chickens.

Disposal of container

Dispose of in a manner consistent with federal, state, and local regulations.