

pharmaniaga®

i-CALCIUM TABLET 300MG

DESCRIPTION

Round, biconvex with scoreline on one side and white in colour.

COMPOSITION

Each tablet contains: Calcium Lactate 300mg providing Calcium 39mg.

PHARMACOKINETICS

Absorption

Calcium is absorbed from the small intestine; about one-third of ingested calcium is absorbed although this can vary depending upon dietary factors and the state of the small intestine. Absorption decreases with age and may be more efficient when the body is deficient in calcium or from diets deficient in calcium, and increases by 1,25-dihydroxycholecalciferol, the active metabolite of Vitamin D. Absorption is increased in calcium deficiency during periods of high physiological requirement such as during childhood or pregnancy and lactation.

Distribution

Calcium is secreted in sweat, skin, hair, nails, saliva, bile and pancreatic juice. Calcium crosses the placenta and there is significant loss of calcium in milk during lactation.

Excretion

Up to 400mg may be excreted in the urine daily; the amount is directly related to the excretion of sodium ions and is also increased by increased dietary magnesium. Urinary excretion of calcium decreases when increased amounts of phosphorus are ingested and increases when the phosphorus intake is reduced.

INDICATIONS

As a health supplement.

RECOMMENDED DOSAGE

Usual Adult Dose : 1 tablet 3 times a day after meal.

ROUTE OF ADMINISTRATION

Oral

CONTRAINDICATIONS

- Severe hypercalcaemia and hypercalciuria (eg hypervitaminosis D, hyperparathyroidism, severe renal failure (must not be given calcium supplements).

WARNINGS AND PRECAUTIONS

- Calcium salts should be administered with caution to patients with impaired renal function or a history of renal stone formation, cardiac disease, or sarcoidosis.

INTERACTIONS WITH OTHER MEDICAMENTS

- Calcium enhances the effects of digitalis on the heart and may precipitate digitalis intoxication.
- Calcium Lactate Tablets must be used with care in patients receiving alternative compound vitamin or mineral preparations, which often contain additional sources of calcium.
- Calcium salts reduce the absorption of tetracyclines.

PREGNANCY AND LACTATION

Pregnancy

There is epidemiological evidence of the safety of calcium in pregnancy.

Lactation

No problems are anticipated with the administration of Calcium Lactate Tablets during lactation

SIDE EFFECTS

Calcium salts may cause constipation.

SYMPTOM AND TREATMENT OF OVERDOSE

Symptom

The symptoms of overdosage with calcium include anorexia, lassitude, nausea, vomiting, headache, extreme thirst, vertigo, raised blood urea, constipation, abdominal pain, muscle weakness, mental disturbance, polydipsia, polyuria; calcium may be deposited in many tissues including the kidney and arteries and the plasma cholesterol level may become elevated. Cardiac arrhythmias and bradycardia may also occur.

Treatment

Calcium intake should be reduced to a minimum and any dehydration and electrolyte imbalance corrected immediately. Severe hypercalcaemia should be treated with an iv infusion of sodium chloride 0.9%; a loop diuretic may be given to increase urinary calcium excretion. If this fails, calcitonin may be administered by injection, or alternative, bisphosphonates, plicamycin or corticosteroids may be used. Phosphate infusion must not be given due to the danger of metastatic calcification. In severe cases, significant amounts of calcium may be removed by peritoneal dialysis.

Patients with symptoms of overdosage should avoid exposure to direct sunlight.

Special care must be exercised when treating overdosage in patients with impaired renal or hepatic function.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINE

None known

PACK SIZE

In box of 30 tablets (3 blisters x 10 tablets)

STORAGE CONDITIONS

Keep in a dry place, below 30°C; Protect from light.

SHELF LIFE

Product should not be used beyond the expiry date imprinted on the product packaging.

REGISTRATION NUMBER

MAL12055053N

KEEP MEDICINES OUT OF REACH OF CHILDREN

JAUHI UBAT-UBATAN DARI KANAK-KANAK

For further information, please consult your doctor or your pharmacist.

Revision Number : 05

Revision Date : 06 Nov 2025

Lt-158.05 Product Registration Holder and Manufacturer

INSPIRAPHARMA SDN. BHD. (200401023395)

LOT 24 & 25, JALAN 8,

KAWASAN PERUSAHAAN BAKAR ARANG,

08000 SUNGAI PETANI, KEDAH DARUL AMAN, MALAYSIA.

145mm

104mm