

FRONT SIDE
SAME SIZE ARTWORK
L : 100mm x H : 220mm

LICSCAB LOTION

Gamma Benzene Hexachloride Lotion USP 1% w/v (LICSCAB Lotion)

Composition:

Gamma Benzene Hexachloride USP1% w/v
Aqueous baseq.s

Description:

A white cutaneous emulsion, with floral fragrance, free from foreign particles.

Pharmacodynamics:

Licscab is a central nervous system (CNS) stimulant when absorbed systematically. Following absorption through the chitinous exoskeleton of arthropods, Licscab is presumed to stimulate the nervous system, resulting in convulsions and death.

Pharmacokinetics:

Absorption

The absorption of Licscab through the skin is highly variable. Although the incidence of adverse effects appears to be higher in infants and young children than in adults, there have been no studies to demonstrate that children absorb more Licscab than adults.

Lotion - In one study, a mean peak blood concentration of 3 nanograms per mL occurred in infants and children 6 hours after total body application of Licscab lotion for scabies.

Distribution

Licscab is highly lipid-soluble and is stored in adipose tissue and other lipophilic tissue. Licscab is also stored in the white matter of the brain and may persist in low levels for up to 2 weeks following topical application.

Metabolism

A study in 9 healthy adults demonstrated a decreased half-life and increased clearance of antipyrine when preceded by a topical Licscab 1% application of 12 hours duration. This suggests that Licscab induces microsomal drug metabolism.

Elimination Half-life

17.9 to 21.4 hours

Indications:

Licscab lotion is indicated for the treatment of scabies infestation caused by *Sarcoptes scabiei*.

Recommended Dose:

For scabies (*SARCOPTES SCABIEI*)

- 1) It is recommended one ounce of Licscab lotion applied in a thin layer to all areas of the body from the neck down and thoroughly washed off after 8 hours. Licscab should not be applied immediately after a bath or shower and should not be used by persons with extensive dermatitis, or in pregnant or lactating women.
- 2) The lotion should be applied thinly and rubbed in thoroughly from the neck down to the soles of the feet, with special attention under nail and to the hands, feet and interdigital areas. Avoid applying Licscab to open cuts or excoriations.
- 3) The mite does not infest areas of sebaceous gland activity in adults or children, and the face and scalp need not be treated. However, scabies may affect the head in infants.
- 4) The lotion should be left on for 8 to 12 hours and removed by thorough washing.
- 5) One application is usually curative although many patients exhibit persistent pruritus after treatment. It is not known how soon after Licscab administration that a second dose can safely be readministered. Retreatment is not recommended.

Route of Administration

Topical

Contraindications:

Gamma Benzene Hexachloride Lotion is contraindicated for premature neonates because their skin may be more permeable than full term infants and their liver enzymes may not be sufficiently developed. It is also contraindicated for patients with known seizure disorders and for individuals with a known sensitivity to the product or any of its components. The other contraindication are Norwegian (crusted) scabies and other skin conditions (e.g, atopic dermatitis, psoriasis) that may increase systemic absorption.

Warning and Precaution:

WARNING - Topical (Lotion)

1. Licscab should only be used in patients who cannot tolerate or have failed first-line treatment with safer medications for the treatment of scabies.
2. Seizures and deaths have been reported following Licscab use with repeat or prolonged application, but also in rare cases following a single application used according to directions. Licscab should be used with caution for infants, children, the elderly, and individuals with other skin conditions (eg, atopic dermatitis, psoriasis) and in those who weigh less than 110 lbs (50 kg) as they may be at risk of serious neurotoxicity.
3. Licscab is contraindicated in premature infants and individuals with known uncontrolled seizure disorders.

- NOT first-line agent
- Use only as directed
- CNS toxicity may be greater in the young; use with extreme caution in infants, children, and pregnant and nursing mothers
- Avoid contact with eyes or mucous membranes
- Avoid inflamed or excoriated skin
- Avoid simultaneous use with other creams, ointments, or oils
- Conditions which may increase the risk of seizure (e.g, HIV, history of head trauma or prior seizure, CNS tumor, excessive alcohol consumption)



BLACK

BACK SIDE
SAME SIZE ARTWORK
L : 100mm x H : 220mm

- Elderly
- Concomitant medications which lower the seizure threshold
- Individuals weighing less than 50 kg (110 lbs)
- Just put a very thin layer of the Licscab Lotion.
- Wash the Licscab Lotion off your skin after 8 to 12 hours.
- You may still itch after you use Licscab Lotion. Even after all the scabies bugs are dead, they can still make your skin itch for a few weeks. This does not mean you need more Licscab Lotion.
- Avoid retreatment with Licscab.

Interaction with Other Medicaments:

Drug interaction: Oils may enhance absorption therefore, simultaneous use of creams, ointments or oils should be avoided.

The following drug interactions and/or related problems have been selected on the basis of their potential clinical significance (possible mechanism in parentheses where appropriate) - not necessarily inclusive (>> = major clinical significance):

Skin, scalp or hair preparation, other, such as creams, lotions, ointments or oils (simultaneous application may increase the percutaneous absorption of Licscab).

Pregnancy and lactation:

Adequate and well-controlled studies in humans have not been done. Because Licscab is absorbed through the skin and has the potential of causing CNS toxicity, some clinicians do not recommend the use of Licscab during pregnancy. If Licscab is used, however, the recommended dosage should not be treated more than twice during pregnancy. Studies in animals have not shown that Licscab causes adverse effects on the fetus.

For breast feeding, problems in human have not been documented; however, Licscab is systematically absorbed and is distributed into breast milk. Although the concentrations found in human blood following topical application of Licscab make it unlikely that breast milk will contain amounts of Licscab sufficient to cause toxicity, an alternate method of feeding the infant should be used for 2 days.

Side effects:

Gamma Benzene Hexachloride is absorbed through the skin and has the potential for CNS toxicity, especially in infants, children and possibly the elderly.

The following side effects have been selected on the basis of their potential clinical significance (possible signs and symptoms in parentheses where appropriate) - not necessarily inclusive.

Those indicating need for medical attention

Incidence rare:

Skin irritation not present before therapy - If Licscab is applied incorrectly or repeatedly, the incidence of skin irritation is increased: skin rash.

Symptoms of CNS toxicity:

Convulsions; dizziness, clumsiness or unsteadiness; fast heartbeat, muscle cramps; nervousness, restlessness or irritability; vomiting.

Those indicating need for medical attention, if they occur and continue or are bothersome after medication is discontinued. Itching of skin - acquired sensitivity to mites and their products may continue for one to several weeks.

Other adverse effects in single patients that have been associated with Licscab are disseminated intravascular coagulation and subsequent death after oral ingestion and aplastic anaemia after prolonged topical exposure (twice daily application for three weeks).

Symptoms and Treatment of Overdose

Symptoms of overdose

Mild to Moderate Toxicity: Vomiting, headache, nausea, dizziness, diarrhea, and tremor.

Severe Toxicity: Seizures, CNS depression, agitation, ataxia, hypotension, respiratory depression, metabolic acidosis, and coma. Children appear to be more susceptible to the toxic effects of Licscab and more likely to manifest severe effects.

For treatment of toxicity:

- If accidental ingestion occurs, it may be life threatening and prompt gastric lavage is recommended.
- Since oils favor absorption, saline cathartics should be administered rather than oily cathartics for intestinal evacuation.
- If CNS manifestation occur, they may be treated by the administration of pentobarbital, phenobarbital or diazepam.

Storage condition:

Store below 30°C. Keep in a cool dry place. Protect from light. Keep bottle tightly closed.

Availability:

Gamma Benzene Hexachloride Lotion is available in Bottles of 100 ml and 1000 ml.

Shelf life:

24 Months

Registration Number: MAL12095096XZ

Manufactured by:

Galentic Pharma (India) Pvt. Ltd.

R-673, MIDC, TTC, Rabale,

Thane Belapur Road,

Navi Mumbai 400 701, India.

Marketing Authorization Holder/Product License Holder:

Unimed Sdn Bhd

No 53 Jalan Tembaga SD 5/2B,

Bandar Sri Damansara,

52200 Kuala Lumpur.

Revision date: Feb 2013

ML416SP1-3



BLACK