

SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm

birth indices and an increase in the number of still-born pups was seen at 80 and 160 mg/kg/day. There are no adequate and well-controlled studies that have been conducted on topically applied Sertaconazole nitrate cream 2%, in pregnant women. Because animal reproduction studies are not always predictive of human response, Sertaconazole nitrate cream 2%, should be used during pregnancy only if clearly needed.

Nursing Mothers:
It is not known if Sertaconazole is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when prescribing Sertaconazole nitrate cream 2%, to a nursing woman.

Route of administration: Topical

SYMPTOMS AND TREATMENT FOR OVERDOSAGE AND ANTIDOTE (S) Overdosage with Sertaconazole nitrate cream 2%, has not been reported to date. Sertaconazole nitrate cream 2%, is intended for topical dermatologic use only. It is not for oral, ophthalmic, or intravaginal use.

PACKING / PACK SIZES
Tube of 20 g

SHELF LIFE
24 months

STORAGE CONDITIONS, USER INSTRUCTIONS AND PHARMACEUTICAL PRECAUTIONS
Store below 30°C. Do not freeze.
Keep all Medicines out of reach of children.
MAL13010002AZ
Date of Revision: 30 January 2024

Manufactured by:



glenmark
PHARMACEUTICALS LTD.
B/2, Mahalakshmi Chambers,
22, Bhulabhai Desai Road,
Mumbai - 400 026, (India).
At: Plot No. E-37, 39, D-Road,
MIDC, Satpur, Nashik - 422 007,
Maharashtra State, India.
© Trade Mark

Product Registration Holder
Glenmark Pharmaceuticals
(Malaysia) Sdn Bhd (660397-W)
D-31-02, Menara Suezcap 1,
No. 2, Jalan Kerinchi,
59200 Kuala Lumpur, Malaysia

PE00000 MY

For the use only of a Registered Medical Practitioner
or a Hospital or a Laboratory

Konzert[™] Cream 2%w/w
Sertaconazole Nitrate Cream 2% w/w

SPACE FOR
PHARMACODE

COMPOSITION
Each gm of cream contains:
Sertaconazole Nitrate BP20 mg
Cream baseq.s.
Preservative: Benzyl Alcohol BP 10 mg

THERAPEUTIC CLASS
D01AC14

DESCRIPTION
A printed carton containing an insert & a lami tube containing white colour semi solid cream. It may contain fine white to off white coloured visible particles.

DOSAGE FORM
Cream

PHARMACOLOGY
Pharmacodynamics:
Mechanism of Action
The mechanism of antifungal action of Sertaconazole nitrate is similar to that of other imidazoles, in that it inhibits fungal biosynthesis of ergosterol, resulting in disorganization of the fungal plasma membrane and altered permeability. Like other imidazoles, sertaconazole has demonstrated a broad spectrum of antifungal activity both in vitro and in experimental in vivo models, which includes dermatophytes (eg, Epidermophyton, Microsporum,

SAME SIZE ARTWORK
LEAFLET SIZE: 140 mm x 88 mm

Trichophyton), opportunistic filamentous fungi (eg, Aspergillus, Alternaria, Scopulariopsis, Fusarium), and pathogenic yeasts such as Malassezia furfur, Candida albicans, Candida tropicalis, Torulopsis, and Trichosporon.

Activity has also been observed against Trichomonas and some gram-positive organisms (staphylococci, streptococci). One preclinical investigation has suggested the greater activity of sertaconazole compared to miconazole (equivalent doses) against dermatophytes and candida. In vitro testing of antifungal agents showed that sertaconazole (MIC 1.36 mg/L) was more active against Candida yeast isolates than bifonazole (MIC 7.2 mg/L) and terbinafine (MIC 13.77 mg/L). Other experimental data suggest the drug may also be more active than other imidazoles.

Pharmacokinetics
Sertaconazole achieves high epidermal concentrations following cutaneous application. Cutaneous absorption was 64% of the dose at 12 hours and 72% at 24 hours following topical application of a 2% cream. Systemic absorption is minimal to undetectable. The drug was undetectable in serum or urine samples from healthy subjects for up to 24 hours with 16 g of sertaconazole 2% cream. In a multiple dose pharmacokinetic study that included 5 male patients with interdigital tinea pedis (range of diseased area, 42 140 cm²; mean, 93 cm²), Sertaconazole nitrate cream 2%, was topically applied every 12 hours for a total of 13 doses to the diseased skin (0.5 grams sertaconazole nitrate per 100 cm²). Sertaconazole concentrations in plasma measured by serial blood sampling for 72 hours after the thirteenth dose were below the limit of quantitation (2.5 ng/mL) of the analytical method used.

INDICATIONS
Sertaconazole nitrate cream 2% is used for the topical treatment of superficial mycoses, such as dermatophytoses, Tinea pedis (athlete's foot), Tinea cruris (Hebra's eczema), Tinea corporis (Herpes circinatus), Tinea barbae (Mycosis of the beard), Tinea Manus, Candidiasis (moniliasis), Pityriasis versicolor (Pityrosporum orbiculare), Malassezia furfur formerly.

CONTRAINDICATIONS
Sertaconazole nitrate cream 2% is contraindicated in patients who have a known or suspected sensitivity to Sertaconazole nitrate or any of its components or to other imidazoles.

SIDE EFFECTS / ADVERSE REACTIONS
In non-US post-marketing surveillance for Sertaconazole nitrate cream 2%, the following cutaneous adverse events were reported: contact dermatitis, erythema, pruritus, vesiculation, desquamation, and hyperpigmentation.

PRECAUTIONS / WARNINGS
Sertaconazole nitrate cream 2%, is not indicated for ophthalmic, oral or intravaginal use.
General: Sertaconazole nitrate cream 2%, is for use on the skin only. If irritation or sensitivity develops with the use of Sertaconazole nitrate cream 2%, treatment should be discontinued and appropriate therapy instituted.

Physicians should exercise caution when prescribing Sertaconazole nitrate cream 2%, to patients known to be sensitive to imidazole antifungals, since cross-reactivity may occur.

Information for Patients: The patient should be instructed to:

1. Use Sertaconazole nitrate cream 2%, as directed by the physician. The hands should be washed after applying the medication to the affected area(s). Avoid contact with the eyes, nose, mouth and other mucous membranes. Sertaconazole nitrate cream 2%, is for external use only.
2. Dry the affected area(s) thoroughly before application, if you wish to use Sertaconazole nitrate cream 2%, after bathing.
3. Use the medication for the full treatment time recommended by the physician, even though symptoms may have improved. Notify the physician if there is no improvement after the end of the prescribed treatment period, or sooner, if the condition worsens.
4. Inform the physician if the area of application shows signs of increased irritation, redness, itching, burning, blistering, swelling or oozing.
5. Avoid the use of occlusive dressings unless otherwise directed by the physician.
6. Do not use this medication for any disorder other than that for which it was prescribed.
7. As this preparation contains benzyl alcohol, its use should be avoided in children under two years of age. Not to be used in neonates.

DRUG INTERACTION
Potential interactions between Sertaconazole nitrate cream 2%, and other drugs or laboratory tests have not been systematically evaluated.

RECOMMENDED DOSAGE, DOSAGE SCHEDULE AND ROUTE OF ADMINISTRATION
The cream is to be applied 1 or 2 times daily (preferably at night, or in the morning and at night), mildly and uniformly on the lesion, trying, besides, to embrace 1cm of sound skin (approximately) around the affected area.
The duration of treatment to obtain healing varies from one patient to another, in function of the etiologic agent and location of the infection. In general, 4 weeks of treatment are recommended to assure a complete clinical and microbiological healing and the non-appearance of relapses, although, in many cases, this clinical-microbiological healing appears earlier, between 2-4 weeks treatment.
Use in the elderly and children: Same as in adults

SPECIAL POPULATIONS
Pregnancy:
Pregnancy Category C
Oral reproduction studies in rats and rabbits did not produce any evidence of maternal toxicity, embryotoxicity or teratogenicity of Sertaconazole nitrate at an oral dose of 160 mg/kg/day (40 times (rats) and 80 times (rabbits) the maximum recommended human dose on a body surface area comparison). In an oral peripostnatal study in rats, a reduction in live

PE00000 MY

ICONGRAPHICS CODE:

PANTONE SHADE



PANTONE
BLACK
PROCESS C

Supersedes
Artwork Code:
PE61201

PHARMACODE: