

PRODUCT NAME: HINABACI PREMIX GRANULAR

COMPOSITION: Each 100g premix contains Bacitracin 15g.

PRODUCT DESCRIPTION: Light brown or tan granule with characteristic odor.

PHARMACODYNAMIC: Bacitracin Zinc has strong anti-bacteria activities to gram-positive bacteria and part of gram-negative bacteria. It has a strong effect to most G+, such as Staphylococcus aureus (including Staphylococcus aureus resistant to Penicillin), streptococcus, streptococcus pneumoniae, enterococcus, et al., also affect to some G- such as meningococcus, influenza bacteria, still has some effect to helicoid and actinomycete. G- bacillus resistant to it.

PHARMACOKINETIC: It can't be absorbed by oral administration, mostly discharged with feces in 2 days. Easily absorbed by intramuscular injection, plasma drug concentration can reach to peak 2 hours after injection. Wide distribution, most in bile, liver, and kidney, little distribute in other tissues, it can be discharged through the kidney quickly. After 24 hours, no residues in other tissues except bile, liver and kidney. By breast injection, residual time in breast no longer than 72 hrs.

INDICATION:

Poultry: For the treatment and prevention of necrotic enteritis

Swine: For aid in treatment of bacterial enteritis (scours) and aid in prevention of bacterial enteritis

USAGE & DOSAGE:

Swine: - As an aid in the treatment of bacterial enteritis (scours) in swine (except coliform diarrhea):
110 mg bacitracin/kg of complete feed and provide this as the sole ration for 5 to 15 days.

- As an aid in the prevention of bacterial enteritis (scours) in swine (except coliform diarrhea):
55 mg bacitracin/kg of complete feed and provide this as the sole ration continuously.

Poultry: - Treatment of necrotic enteritis: 200g bacitracin activity per tonne to be fed for 5-7 days

- Aid in prevention of necrotic enteritis: 40g bacitracin activity per tonne. Medicated feed should be
Fed to poultry during the period when they are at most risk of the disease, which is usually between
2 and 12 weeks.

CONTRAINDICATIONS: None Known

WARNING & PRECAUTIONS: Pigs refusing to eat should be treated individually.

DRUGS INTERACTIONS:

Bacitracin zinc could combine use with penicillin, streptomycin, neomycin, polymyxin.

Do not use in feed containing bentonite or other pellet binding agent.

PREGNANCY AND LACTATION: Can not be used in the laying period.

ADVERSE EFFECTS: None Known

OVERDOSE AND TREATMENT: Stop medication and to seek for veterinarian advice if there is sign of
overdosage.

WITHDRAWAL PERIODS: 0 days before slaughter.

STORAGE: Keep sealed and stored in cool and dry place below 25°C.

PACKAGING AVAILABLE: 25 kg packed into Polyethylene plastic bag.

SHELF LIFE: 2 YEARS

MANUFACTURE BY:

HINAPHARM PHARMACEUTICAL CO., LTD. FOSHAN

Add: Shashuihe West Road , Cangjiang Industrial Zone , Gaoming District , Foshan ,Guangdong, China

MARKETING AUTHORISATION HOLDER:

PAHANG PHARMACY SDN. BHD.

Add: Lot 5979, Jalan Teratai, 5 ½ Mile Off Jalan Meru,
41050 Klang, Selangor Darul Ehsan, Malaysia.

DATE OF REVISION OF PACKAGE INSERT: May 26, 2015