

CD.JEVAX® Inj

Japanese Encephalitis Vaccine, Live

Description

CD. JEVAX®, Japanese Encephalitis Vaccine, live is a sterile, lyophilized vaccine for subcutaneous use, prepared by passaging Japanese Encephalitis (JE) virus, strain SA14-14-2 in monolayer of primary hamster kidney cell culture.

The product looks like a light yellow crisp cake. After reconstitution, it shall turn into a clear, orange-red or light pink liquid.

Ingredients

A human dose of 0.5mL contains: -

Live-attenuated JE virus (Seed strain: SA14-14-2)	5.4 log PFU above
Gelatin	4.8mg less
Sucrose	21mg less
Lactose	21mg less
Urea	2.4mg less
Human Serum Albumin	3.0mg less
Sterile Diluent (Sterile water for injection)	0.5mL

Indication and Usage

Prophylactic immunization against Japanese Encephalitis.

Dosage and Administration

-Primary immunization for persons at least 9 months of age and older, a single dose of 0.5mL subcutaneously and a booster dose can be given from 3 months to 1 year after primary dose.

-After reconstitution with the diluent (Sterile water for injection), the vaccine should be used within 1 hour.

- The deltoid muscle of right or left arm is the recommended site of injection. The skin at the site of injection first should be disinfected with alcohol or iodine tincture and allowed to dry thoroughly before injection.

- The vaccine should not be injected into a blood vessel.

- Vaccine recipients should be monitored during and after receiving CD.JEVAX® and if any reactions (fever, convulsion etc.) are observed a healthcare provider should be contacted.

- As for any vaccine, adequate treatment provision, including epinephrine, should be available for immediate use in case an anaphylactic reaction occurs.

Warning and Precautions

- Prior to injection the health care provider should question the vaccinee or his/her parent, or guardian on his/her recent medical history.
- Remove plastic tab of flip-off cap. Do not remove the rubber stopper. Cleanse stopper with a suitable disinfectant. Reconstitute only with the diluent (water for injection). Shake vial thoroughly.
- The vaccine should be inspected visually for extraneous particulate matter and/or discoloration prior to administration whenever solution and container permit. If either of these conditions exists, the vaccine should not be administered.
- The vaccine should be reconstituted just before use and not be frozen or stored again after reconstitution.

Pregnancy and Lactation

- Do not administer CD. JEVAX® during pregnancy or lactation.

Contraindications

- Persons with a proven or suspected history of hypersensitivity/ anaphylactic reaction to any component of the vaccine, including gelatin
- Persons with fever, acute infectious disease, tympanites or active untreated tuberculosis
- Persons with malnutrition, general allergy and convulsion
- Persons with cardiac, liver or kidney troubles
- During pregnancy or lactation
- Persons undergoing any type of immunosuppressive therapy
- Persons with a weak or not proper functioning immune system

Drug Interactions

- Although no interactions are known with other medication, the healthcare provider should question the vaccinee, parent or guardian on recent or current medication usage.
- Currently no clinical data is available on the administration of CD. JEVAX® given at the same time with any other vaccine. However, it is advisable to observe a period of at least 2 to 4 weeks when given prior or after to CD. JEVAX®.

Adverse Reactions

- As is the case for all medications, the administration of CD. JEVAX® can also cause adverse reactions.
- Clinical reactions are observed in a small percentage of the vaccinees after administration of CD. JEVAX®.
- Some events listed as below have been reported after injection, which normally does not last longer than 2 days, mostly relieved spontaneously; commonly no particular treatment is required, in case of necessity, symptomatic treatment might be recommended.
 - Fever (Increase temperature above 37.5°C)
 - Rash and Nausea
 - Local redness, pain or sensitivity
 - Crying, loss of appetite, sleepiness or sleep problems (children)

Pack Size

Vial, single dose (10 per package) with/without vial or ampoule of diluents (10 per package, each 0.5mL)

Storage

The vaccine should be stored and shipped at 2-8°C, protected from light. The diluent should be stored at between 2-30°C.

Expiry Date

The vaccine shall be used before the expiry date (Exp. Date) as indicated on the label.

Manufacturer

Chengdu Institute of Biological Products Co. Ltd,
Chengdu, P.R. China

Worldwide Distributor

Glovax Co., Ltd, Seoul, South Korea

Registration No.: MAL13025032AZ

Product Registration Holder:

Farmaco Healthcare Sdn. Bhd.
E-05-09, Sunway Geo Avenue, Jalan Lagoon Selatan,
Bandar Sunway, 47500 Subang Jaya, Selangor, Malaysia.

Revision: Jan 2019
PI0401_E1505_rev.03.