

Product Name:

FLORCOSOL 10% Oral Solution

Composition:

Each ml contains:

Florfenicol 100mg

Product Description:

Light yellow solution

Pharmacodynamics:

Swine: Florfenicol is a synthetic broad spectrum antibiotic active against many gram-negative and gram-positive bacteria isolated from domestic animals. It acts by inhibiting bacterial protein synthesis. Swine: In vivo activity has been demonstrated against bacterial pathogens commonly associated with swine respiratory disease, including *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*. In vitro activity has been demonstrated against *Salmonella* spp and *Streptococcus suis* and in vitro activity against *Mycoplasma hyopneumoniae* and *Mycoplasma hyorhinis* has been detected. However, the clinical significance of this activity in swine is not known.

Pharmacokinetics:

After administration to pigs by gavage at 10mg/kg under experimental conditions, absorption of florfenicol was variable but peak serum concentrations of approximately 5microgram/ml were reached approximately 3 hours after dosing. The terminal half-life was between 3 to 4 hours. After a single treatment the serum concentrations of florfenicol remained above 1 microgram/ml for 12 to 18 hours.

After absorption and distribution, florfenicol is extensively metabolised by pigs and rapidly eliminated, primarily in urine and the faeces in a ratio of 3:1

A fraction is secreted unchanged, the rest is metabolised to 5 metabolites

Indications:

SWINE: Indicated for the treatment of swine respiratory disease susceptible to florfenicol associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

Dosage and Administration:

1.0ml of Florcosol 10% Oral Solution to 1 litre drinking water corresponds to a 100mg/L drinking water concentration. Swine: 100 ppm (100 mg/L) for 5 consecutive days in drinking water. This dosage rate is equivalent to 10 mg/kg body weight for pigs that are drinking 10% of their body weight.

Contraindications:

Do not use in boars intended for breeding purpose.

Do not administer in cases of previous allergic reactions to florfenicol.

Interaction with Other Medicaments:

None known

Pregnancy and Lactation:

The safety of the product during pregnancy and lactation has not been studied in sows. Use of the product during pregnancy and lactation is therefore not recommended. Do not use in breeding boars because toxicity studies in rats have shown adverse effects on the male reproductive system.

Side effects:

Commonly observed adverse effects are diarrhoea, perianal inflammation and rectal eversion. These effects are transient, resolving on cessation of treatment. Increased serum calcium may also be observed.

Symptoms and Treatment of Overdose:

In the event of overdose, a reduction in food and water consumption, together with a decrease in bodyweight may be observed. There may be an increase in refused feed and an increase in serum calcium.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Shake well before use. The treated pigs should be placed under special observation. On each of the five days of treatment, untreated food should not be given until the full daily amount of medicated feed has been ingested by the pigs. If there is no significant improvement after 3 treatment days, the diagnosis should be reviewed and if necessary the treatment should be changed. Treatment should not exceed 5 days.

Withdrawal Periods:

For meat

Swine: 21days

Broiler chickens: 7days

Container disposal: Dispose of any unused product and empty container in accordance with local waste regulation authority.

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,
Prai Industrial Estate MK1,
13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

DEXCHEM INDUSTRIES SDN BHD

No. 6 & 8, Jalan Industri Cherok To'Kun 2,
Taman Industri Cherok To'Kun,
14000 Bukit Mertajam, Pulau Pinang.

DATE OF REVISION: APRIL 2019