

VETFLOR 25%w/v Oral Solution

FOR ANIMAL USE ONLY

Product Name :

VETFLOR 25%w/v Oral Solution.

Name and Strength of Active Substance(s)

Each ml contains 250 mg florfenicol and 10 mg benzyl alcohol (as preservative).

Product Description

VETFLOR 25%w/v Oral Solution is a yellowish to brownish liquid.

Pharmacodynamics

Florfenicol is a synthetic broad spectrum antibiotic active against many gram- negative and gram- positive bacteria isolated from domestic animals. It acts by inhibiting bacterial protein synthesis.

Pharmacokinetics

After the first oral gavage administration of florfenicol of 42mg/kg bw/day for 3 days to broiler chicken, the highest concentration of florfenicol in serum was measured after 30 minutes then declined to reach 810 µg/ml, 8 hours later. Florfenicol could not be detected in the serum, 60 hours after the last administration. During a continuous treatment of broiler chicken with florfenicol in drinking water at a daily dose of 26mg/kg bw for 18 hours for 3 days, the estimated mean serum concentration of florfenicol was 711 µg/ml. No florfenicol was detected in the serum, 72 hours after the terminal dose. After repeated oral administration of ¹⁴C-florfenicol, given twice daily 12 hours apart at 20mg/kg bw/dose for three days, 93.7% and 98.2% of total radioactivity administered were excreted within 1 day and 7 days after the last dose. In excreta, at 7 days, the parent compound represented the major fraction of the radioactivity (42%), florfenicol amine (25%), florfenicol oxamic acid (5%) and florfenicol alcohol (10%) and the remaining part of radioactivity being represented by a small percentage of three unknown compounds. No monochloroflorfenicol was detected in excreta. In a radiometric study, chicken received ¹⁴C-florfenicol by oral gavage twice daily 12 hour apart at 20 mg/kg bw/dose for three days (i.e. 40 mg/kg bw/day for three days). Twenty-four hour after the end of oral administration, the levels of radioactivity in edible tissues were 146 µg equivalents florfenicol/kg in muscle, 475 µg/kg in skin+ fat, 11148 µg/kg in liver and 3125 µg/kg in kidney. Then declined to reach approximately 50 µg/kg in muscle and in skin + fat 5 days after the end of treatment, whereas in liver and kidney significant amounts of residues were still found : 2403 and 844 µg equivalents florfenicol/kg respectively. Florfenicol amine was the major metabolite measured in edible tissues of chicken, florfenicol, the only microbiologically active compounds, being only detected in skin + fat in low concentrations (4%) and in kidney (1%). In a non-radiometric study, broiler chickens received florfenicol according to the recommended regimen via drinking water at concentrations equivalent to 17 to 30 mg/kg bw/day for 3 days. At 0.5 days after the end of the treatment the concentrations of florfenicol amine were lower than the limit of quantification for muscle (less than 50 µg/kg) and for skin + fat (less than 109 µg/kg) whereas in liver and kidney, significant amounts of florfenicol-amine were measured 2862 and 1161 µg/kg respectively. Florfenicol amine concentrations declined to reach 2038 and 679 µg/kg in liver and kidney, 1 day after the end of the treatment and to 1215 and 484 µg/kg in liver and kidney, 3 days after the end of the treatment. Seven days after the end of the treatment, the concentrations of florfenicol amine were below the limit of quantification for liver (less than 461 µg/kg) and could be still measured in kidney (136 µg/kg).

Indication

Broiler chicken: For the treatment and control of air sacculitis associated with E. coli susceptible to florfenicol.

Recommended Dosage

100ppm (100mg florfenicol/ L) for 3 consecutive days in drinking water. This dosage rate is equivalent to 35mg florfenicol/kg bodyweight of birds. Solutions with concentrations comprised between 1.2g and 12g of florfenicol per liter precipitate. If proportioner is used, the proportioner should be tested for accuracy before use.

Direction of Use :

To be administered via animal drinking water. Mixing Directions : The liquid concentrate must be added in the amount of animal drinking water estimated to be consumed within 12 – 24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water must be prepared each day.

Contraindications

The use of this drug is contraindicated in animals with a history of an allergic reaction to florfenicol. Do not feed to birds intended for breeding purposes.

Warnings and Precautions

The medicated water should be administered as the only source of drinking water for 3 consecutive days. Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol. To limit the development of antimicrobial resistance, the choice of this drug as the most appropriate treatment should be confirmed by clinical experience supported where possible by pathogen culture and drug susceptibility testing. Not for human use. Keep out of reach of children. Avoid direct contact with skin, eyes and clothes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. Consult a physician if irritation persists.

Interactions with Other Medicaments

There is inadequate data on adverse drug interactions produced by florfenicol.

Statement on usage during pregnancy and lactation

The effects of florfenicol on reproductive function of treated chickens have not been determined.

Adverse Effects / Undesirable Effects

None known.

Overdose and Treatment

None known.

Withdrawal Period(s)

5 Days

Shelf-life

2 years from date of manufacture. Opened bottle and unused medicated water to be discarded after 24 hours.

Maximum Residual Limit (MRL)

Chicken: Muscle 100ug/kg, Skin & Fat: 200 ug/kg, Liver: 2500ug/kg, Kidney: 750ug/kg.

Storage Conditions

Store between 15 - 25°C. Keep medicine out of reach of children. Jauhkan ubat dari kanak-kanak.

Disposal of Containers

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned empty bottle can be safely disposed according to local government regulations.

Dosage Forms and packaging available

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2013-06-21