

YSP TREN

INJECTION 100MG/ML

Ingredient(s):

Each ml contains:

Tranexamic Acid100mg

Pharmacology (Summary of Pharmacodynamic and Pharmacokinetic):**Pharmacodynamic**

Tranexamic acid is an anti-fibrinolytic agent. It acts principally by inhibiting plasminogen activators which have fibrinolytic properties and to a lesser degree by inhibiting plasmin activity.

Pharmacokinetic

1. Tranexamic acid has the action and uses of aminocaproic acid but is reported to be about 10 times more potent. It is useful in enhancing hemostasis when fibrinolysis contributes to bleeding.
2. Tranexamic acid is rapidly absorbed from the gastro-intestinal tract following oral administration. Up to 40% of a dose by mouth and 90% of a dose by intravenous injection is excreted in the urine within 24 hours. It diffuses across the placenta.
3. The serum half-life after intravenous was considered to be 1 to 3 hours.

Indication(s):

1. Bleeding tendencies in which systemic hyperfibrinolysis is considered to be involved. (Leukemia, aplastic anemia, purpura etc., abnormal bleeding during or after operation.)
2. Abnormal bleeding in which local hyperfibrinolysis is considered to be involved. (pulmonary hemorrhage, nasal hemorrhage, vaginal hemorrhage, renal hemorrhage, abnormal bleeding during or after prostate surgery.)
3. Symptoms, such as erythema, swelling or pruritus in the following diseases: Eczema or similar conditions, urticaria, drug eruptions or toxicoderma
4. Symptoms, such as pharyngalgia, redness, hyperemia or swelling in the following diseases: Tonsillitis, pharyngolaryngitis.
5. Pain in the oral cavity or mucosal aphtha in cases of stomatitis.

Dosage and Administration:

For intramuscular or intravenous injection, 250-500mg daily in divided doses.

Mode of Administration:

Intramuscular or intravenous injection

Contraindication(s):

Tranexamic acid should not be used in the presence of disseminated intravascular coagulation (DIC) without concomitant heparin.

Precaution(s) / Warning(s):

1. Tranexamic acid should be given in reduced doses to patient with impaired renal function.
2. Its use is not advised in patients with intravascular coagulation and not recommended for bleeding into cavities.
3. The risk of clotting may be increased in patients taking oral contraceptives and in those with underlying thrombotic states.
4. The preparation should not be used in patients with acquired defective colour vision. If a colour vision disorder should occur during the course of treatment, the drug should be discontinued.

WARNING:

This preparation contains Sodium metabisulphite that may cause serious allergic type reactions in certain susceptible patients. Do not use if known to be hypersensitive to bisulphites.

Interaction with Other Medicaments:

None has been mentioned.

Pregnancy and Lactation:**Pregnancy**

Tranexamic acid does cross the placenta however there are no adequate and well-controlled studies of tranexamic acid use during pregnancy. The usual caution with the use of drugs in pregnancy should be observed.

Lactation

There are limited data on nursing an infant while on tranexamic acid. Data have shown that only minimal amounts of the drug are excreted in breast milk. Levels of tranexamic acid found in breast milk are about one hundredth of those in the maternal serum. Tranexamic acid should only be used in a nursing mother only if clearly needed and use caution when tranexamic acid is administered to a nursing mother.

Side Effect(s):

Tranexamic acid appears to be well tolerated. It can produce dose-related gastrointestinal disturbances. Hypotension and dizziness have occurred, particularly after rapid intravenous dosage. Thrombotic complications (including cerebral thrombosis and central retinal venous and arterial occlusion) have been reported in patients receiving tranexamic acid, but these are usually a consequence of its inappropriate use. Ocular and visual disturbances, including disturbance of colour vision, have been associated with use of tranexamic acid; in such cases the drug should be stopped. Other adverse effects that have been reported include hypersensitivity reactions, skin reactions, musculoskeletal pain, and convulsions.

Symptoms and Treatment for Overdosage, and Antidote(s):

There is no specific antidote for its overdose. Therefore, management of the patient consists of symptomatic and supportive therapy.

Incompatibilities:

Tranexamic Acid should not be added to blood for transfusion, or to injections containing penicillin.

Shelf-Life:

3 years from the date of manufacture.

Storage Condition(s):

Store at temperature below 30°C.

Product Description:

Solution in the container: A clear and colorless solution.

Solution after mixing with compatible solution: A clear and colorless solution.

Dosage Forms and Packaging available:

5ml x 10, 5ml x 25 & 5ml x 50 vials.

5ml x 10, 5ml x 50 & 5ml x 100 clear ampoules.



Manufacturer and Product Registration Holder:

Y.S.P. INDUSTRIES (M) SDN. BHD. (199001001034)

Lot 3, 5, 7, 12 & 14, Jalan P/7, Section 13,
Kawasan Perindustrian Bandar Baru Bangi,
43000 Kajang, Selangor, Malaysia.

Ordering Line: 1 800 88 3027

Product Info: 1 800 88 3679