

Beutlich®

PHARMACEUTICALS LLC

Since 1954

HurriCaine Cherry Topical Anesthetic Gel 20%

Amber, opaque gel with a cherry odor.

Content:

Benzocaine 20%

Indication:

To be applied topically to oral tissues for temporary relief of pain associated with dental conditions.

Pharmacology:

Benzocaine is a topical anesthetic. It is an analgesic (dulls pain), anesthetic (numbs nerves), antipruritic (calms itching). It works on the mucosal tissue, because the nerve endings are closer to the surface.

Pharmacokinetics:

Temporary contact with oral mucosa only. No specific human systemic absorption studies have been performed. Benzocaine is not known to be absorbed through the oral mucosa in amounts which would be considered harmful.

Contraindications:

Patients with known hypersensitivity to benzocaine. Application to severely traumatized mucosa or to areas of sepsis.

Side effects/Adverse reactions:

Do not use if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. Do not use if imprinted seal under cap is broken or missing. Contact dermatitis and/or hypersensitivity to benzocaine can cause burning, stinging, pruritus, tenderness, erythema, rash, urticaria and edema. Rarely, benzocaine may induce methemoglobinemia causing respiratory distress and cyanosis. Methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. Stop use and seek immediate medical attention if you or a child in your care develops: • weakness • confusion • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy • pale, gray, or blue colored skin (cyanosis). Symptoms may appear up to 2 hours after use even if you have used this product before.

Precautions:

Localized allergic reactions may occur in patients sensitive to benzocaine or after prolonged or repeated use. Cross-sensitivity may occur with other local anesthetic of the ester type and systemic medications such as sulphonamides, thiazides and paraamino dyes. Methemoglobinemia may rarely occur, particularly in infant and children, following application of large amounts e.g. in pharyngeal anesthesia. Topical benzocaine should not be used for teething or in children under 2 years of age. Children under 12 years of age should be supervised in the use of this product. It should not be used for prolonged periods.

Use in pregnancy and lactation:

Animal reproduction studies have not been conducted with benzocaine. Topical gel should be given to a pregnant woman only if clearly indicated. It is not known whether benzocaine is excreted in human milk. Caution should be exercised when topical gel is administered to a nursing woman.

Drug Interactions:

None known.

Route of Administration:

Oral administration only

Dosage and administration:

Apply small amount evenly to the affected area as directed by the physician.

Effects on Ability to Drive and Use Machine:

Active substance benzocaine has negligible influence on the ability to drive and use machines. See also Precautions section.

Treatment of overdose:

Accidental swallowing of large quantities causes numbness, nausea, vomiting, irritation to eyes. In case of eye and skin contact, flush with large quantities of water. In case of accidental swallowing, to get medical attention immediately. Treatment is primarily supportive. It includes monitoring for development of methemoglobinemia.

Storage conditions:

Store in a cool, dry place below 30 °C. Protect from light. Keep out of reach of children.

Presentation : 30gm/bottle

Shelflife : 2 years

Manufactured by : Dental Technologies Inc.
6901 N, Hamlin Avenue,
Lincolnwood, IL 60712
United States

Distributed by : TK Dental Trading
488B-03-24, Midlands Park,
Jalan Burmah,
10350 Georgetown,
Penang, Malaysia

Date of revision: 04-18-2021