

TILCOSIN 250 Oral Solution

FOR ANIMAL USE ONLY

Brand or Product Name :

TILCOSIN 250 Oral Solution

Name and Strength of Active Substance(s)

Each ml contains Tilimicosin Phosphate 250mg (equivalent to 224.7mg Tilimicosin) and 0.1 %w/v Potassium Sorbate as preservative.

Product Description

TILCOSIN 250 Oral Solution is a light yellowish to brownish liquid.

Pharmacodynamics

Tilimicosin is a semi-synthetic antibiotic of the macrolide group and is believed to affect protein synthesis. It has bacteriostatic action but at high concentrations it may be bactericidal. This antibacterial activity is predominantly against Gram-positive microorganism with activity against certain gram-negative ones and *Mycoplasma* of a bovine, porcine, ovine and avian origin. In particular, its activity has been demonstrated against the following microorganism : Chickens and turkeys *Mycoplasma gallisepticum* and *Mycoplasma synoviae*. Scientific evidence suggests that macrolide act synergistically with the host immune system. Macrolides appear to enhance phagocyte killing of bacteria. Tilimicosin has been shown to inhibit in vitro the replication of the Porcine Reproductive and Respiratory Syndrome virus in alveolar macrophages in a dose dependent fashion. Cross-resistance between tilimicosin and other macrolides and lincomycin has been observed.

Pharmacokinetics

Poultry: As early as 6 hours after oral administration of 75 mg tilimicosin/l drinking water, the average active substance concentrations detected in lung and alveolar tissue were 0.63 µg/g and 0.30 µg/g respectively. 48 hours after the start of treatment, the tilimicosin concentrations in lung and alveolar tissue were 2.3 µg/g and 3.29 µg/g respectively.

Indication

Chickens : For the treatment and prevention of respiratory disease in chickens flocks, associated with *Mycoplasma gallisepticum* and *M. synoviae*.

Recommended Dosage

Chickens (except hens producing eggs for human consumption): To be included in the drinking water at a daily dose of 15-20 mg/kg bodyweight in chickens for 3 days, which may be achieved by the inclusion of 75 mg tilimicosin per litre (1 Litre Tilcosin 250 per 3000 Litres drinking water).

Mode of Administration

To be diluted in the animal drinking water. Medicated drinking water should be prepared fresh every 24 hours, using only clean water.

Contraindications

Do not allow horses and other equines access to drinking water containing tilimicosin.

Do not use in case of hypersensitivity to tilimicosin or to any of the excipients.

Warnings and Precautions

Special warnings for each target species

Important: Must be diluted before administration to animals. Chickens: Water consumption should be monitored in order to guarantee adequate dosing. In case water consumption does not match quantities for which recommended concentrations were calculated, the concentration of Tilcosin 250 has to be adapted in a way that the recommended dosage will be taken up by the animals or different medication should be considered.

Special precautions for use in animals

For oral use only. Inappropriate use of the product may increase the prevalence of bacteria resistant to tilimicosin and may decrease the effectiveness of treatment with tilimicosin-related substances. The use of the product should be based on susceptibility tests. Medicated drinking water should be prepared fresh every 24 hours.

Special precautions to be taken by the person administering the veterinary medicinal product to animals

Tilimicosin may induce irritation. Macrolides, such as tilimicosin, may also cause hypersensitivity (allergy) following inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilimicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves, do not eat, drink or smoke when handling this product. Wash hands after use. In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Do not handle the product if you are allergic to ingredients in the product. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention.

Interactions with Other Medicaments

None Known.

Usage during pregnancy and lactation

The safety of tilimicosin has not been established in animals used for breeding purposes.

Adverse Effects

In very rare cases, a decrease in water intake has been observed.

Overdose and Treatment

No symptoms of overdose have been seen in chickens given drinking water containing levels of tilimicosin up to 375mg/litre (equivalent to 75-100 mg/kg bodyweight or 5 times the recommended dose) for 5 days, daily treatment with 75 mg/litre (equivalent to the maximum recommended dose) for 10 days resulted in a reduction in faecal consistency.

Withdrawal Period(s)

Chickens : 12 days. Not authorized for use in laying birds producing eggs for human consumption. Do not use within 14 days of onset of the laying.

Shelf-life

2 years from date of manufacture. Opened bottle must be discarded after 24 hours. Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL)

Muscle, fat (cattle, poultry, pig and sheep) = 100 µg /kg , [Ref. REGOV, Version 2, dated Dec 2009], Liver (chicken) = 1000µg /kg, kidney (chicken) = 250µg /kg.

[Ref.EMA/MRL/764/00-FINAL, dated Dec 2000]

Storage Conditions

Store in below 25°C. Keep medicine away from children. Jauhkan ubat dari kanak-kanak.

Disposal of containers

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned empty bottle can be safely disposed in trash according to local government regulations.

Packaging available

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2013-11-11