



Cefasily[®]

Film coated tablets

Read all of this leaflet carefully because it contains important information for you.

This medicine is available without prescription.

- Nevertheless you still need to use **Cefasily[®]** film coated tablets carefully to get the best results from it.
- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- You must see a doctor if your symptoms worsen or do not improve.

In this leaflet:

1. What **Cefasily[®]** film coated tablets is and what it is used for
2. Before you take **Cefasily[®]** film coated tablets
3. How to take **Cefasily[®]** film coated tablets
4. Possible side effects
5. Storing **Cefasily[®]** film coated tablets
6. Further information

Composition:

Each 425 mg film coated tablet contains:

Active ingredient:

Dry extract from *Silybum marianum* fruits (35-45:1) 200 mg
corresponding to 52.5 % Silymarin (calc. as Silibinin, HPLC)

Extraction solvent: Acetone

Excipients:

Microcrystalline cellulose, lactose monohydrate (BSE free bovine source), colloidal anhydrous silica, copovidone, magnesium stearate, croscopovidone, povidone, macrogol, hypromellose, talc, titanium dioxide (E 171), ferric oxide (E 172), polyacrylate.

1. WHAT CEFASILY[®] FILM COATED TABLETS IS AND WHAT IT IS USED FOR

Film coated tablets for intake.

Folding boxes with 3 blisters of 20 film coated tablets each.

This is a traditional medicine.

Traditionally used as a liver tonic.

2. BEFORE YOU TAKE CEFASILY[®] FILM COATED TABLETS

Contraindications: This medicine has not been researched for children's use. Please refrain from giving the tablets to children under 12 years of age. There has not been sufficient research for the application of this medicine during pregnancy and lactation. Therefore it should not be used during this period.

Appropriate precautions for use; special warnings: Not applicable.

Please consult your pharmacist/doctor before taking this product. Keep out of reach of children. Jauhi dari kanak kanak.

Interaction with other medicinal products: None known.

>>

3. HOW TO TAKE CEFASILY® FILM COATED TABLETS

Dosage: Unless otherwise prescribed: Adults and children over 12 years take 1 film coated tablet 3 times a day.

Please consult your pharmacist/doctor before taking this product.

Method and/or route(s) of administration: Cefasily® film coated tablets can easily be swallowed with some liquid like water after the meals.

Duration of treatment: At longer persistent complaints, you should consult your doctor.

If you forget to take Cefasily® film coated tablets: Continue as usual, but do not take a double dose to make up for forgotten individual dose.

Effects when treatment with Cefasily® film coated tablets is stopped: None.

4. POSSIBLE SIDE EFFECTS

A mild laxative effect has been observed in very few cases.

If you notice any side effects not mentioned in this leaflet, please inform your physician or pharmacist.

5. STORING CEFASILY® FILM COATED TABLETS

Store below 30° C. Protect from light and moisture.

Do not use after the expiry date stated on the blister and carton.

6. FURTHER INFORMATION

For any information about this medicinal product, please contact the local representative of the Marketing Authorisation Holder.

Product Holder:

Bio-Pharmaceuticals Sdn Bhd
48-2, Jalan Sungai Burong AA32/AA, Sek 32,
40460 Bukit Rimau,
Shah Alam, Selangor
Malaysia

Tel.: + (603) 51319343/42, Fax: + (603) 51319344

email: karen@biopharmaceuticals.com.my

Manufacturer and product owner:

Cefak KG,
Ostbahnhofstr. 15,
D-87437 Kempten,
Germany

Last updated: **June 2017**