

Change of product name from Imequyl 50FG
Flumequine 50%w/w Oral Powder to
Rhomiquyl 50FG Oral Powder 50%w/w

RHOMIQUYL® 50FG Oral Powder 50%w/w

Composition: Flumequine 50 %w/w (as flumequine potency)

Product Description: White to off-white powder

Pharmacodynamics: Flumequine is active against Gram (-) bacteria. Flumequine is a synthetic antibacterial agent which belongs to the Quinolone family. It is the first fluorinated quinolone, with a fluorinate atom which enhances its intrinsic activity and antibacterial spectrum. It acts on nucleic acids by stopping DNA replication of the bacteria. The exact molecular target is the DNA gyrase enzyme involved in the DNA spiral formation. Antibacterial activity of Flumequine is the result of the combination of bacterial cell penetration and DNA gyrase-inhibiting activities.

Pharmacokinetics:
Absorption and distribution
Flumequine is well absorbed. Its bio-availability factor approximates 70 - 75%. Tmax is reached after more or less 13 - 14 h and Cmax varies from 1.7 to 3.7 mg/l. Pharmacokinetic parameters are linearly related to the administered dose of flumequine. No accumulation in blood and no enzymatic induction or inhibition is noticed.

Biotransformation
Flumequine is essentially metabolised into 7-hydroxyflumequine in the kidney.

Elimination
Flumequine has a relatively slow elimination phase. Its half-life approximates 5 - 6 hours. Flumequine is mainly excreted via urine.

Indication:
In cattle, sheep and goats: Curative and preventive treatment in:
- digestive tract infection caused by *Escherichia coli* and *Salmonella spp.*
- respiratory tract infection due to *Pasteurella multocida* and *Haemolytical manheimia*.
In pigs: Curative and preventive treatment in:
- digestive tract infection caused by *Escherichia coli* and *Salmonella spp.*
- respiratory tract infection due to *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *Haemophilus parasuis*.
- urinary tract infection due to *E. coli*.
In poultry: Curative and preventive treatment in:
- respiratory tract, digestive tract and septicemic infections caused by *Escherichia coli*, *Salmonella spp.* and *Pasteurella spp.*

Routes of administration: Oral. To be included in animal feed

Dosage:
To be included in animal feed.
In cattle, sheep and goats: 12mg flumequine per kg body weight per day for 5 days orally.
In pigs: 12mg flumequine per kg body weight per day for 3 to 5 days orally.
In poultry: 12mg flumequine per kg body weight per day for 5 days orally.

Contraindications: Do not use in case of hypersensitivity to quinolones.

Warnings and Precautions:
Special warnings for each target species: None.

Special precautions for use in animals:
The use of quinolones should be based on susceptibility testing whenever possible, and take into account official and local policies for the use of antibiotics. The use outside the recommendations of the SPC may increase the prevalence of bacteria resistant to quinolones and may decrease the effectiveness of treatment with other (fluoro) quinolones due to the potential for cross resistance.

Special precautions to be taken by the person administering the veterinary medicinal product to animals
Handle this product with care and avoid direct contact. Take all recommended precautions by wearing protective clothing, glasses, gloves and mask. Avoid contact with skin and eyes. In case of symptoms such as cutaneous eruption, consult a doctor and show him this information. Oedemas at the level of the face, the eyelids, the lips or respiratory difficulties constitute severe symptoms and require urgent medical cares. Avoid handling this product or the supplemented water in case of allergy to the active substances.

Special precautions for the disposal of unused veterinary medicinal product or waste materials derived from the use of such products
Any unused veterinary medicinal product or waste materials derived from such veterinary medicinal products should be disposed of in accordance with local requirements.

Interactions with other medicaments: Do not associate with trimethoprim -sulphonamide combination.

Statement on usage during pregnancy and lactation:
No embryotoxic, teratogenic or reproduction were observed in studies with laboratory animals. The safety of flumequine was not studied in the target species. The use of specialty during pregnancy and lactation should be an evaluation of the benefit / risk assessment by the veterinarian.

Adverse Effects: None known.

Overdose and Treatment: None known.

Storage Conditions: Store in dry place below 30°C. Protect from direct sunlight.

Withdrawal Period:
Meat and offal: 2 days.
Milk: 2 days.
Eggs: In the absence of waiting time for eggs, do not use in producing species eggs for consumption, four weeks before the start of laying and during it.

Packing: 25 kg
FOR VETERINARY USE ONLY

Registration holder and Manufactured by:

RHONEMA MALAYSIA SDN BHD (200001023279)
Lot 18A & 18B, Jalan 241, Seksyen 51A, 46100 Petaling Jaya, Selangor, Malaysia.
Tel: 03-7873 7355 Fax: 03-2770 0119
E-mail: customercareline@rhonema.com

Date of revision: November 2019
5LRRHOMF3-25K 01/NOV19

Correction of
company address

Change of fax
number

LABEL CODE:		VERSION	COLORS
5LRRHOMF3-25K			 Black
PRODUCT: RHOMIQUYL 50FG ORAL POWDER 50%W/W		Version: 1	
PACK SIZE: 25 KG		MMM/YY: NOV 19	
COUNTRY: MY			
DIMENSIONS: cm = 10 x 14			! Please use the official PANTONE® (P) matching • system for an accurate color representation.

Marketing approval:	Regulatory approval:	GMP approval:
COMPLETE NAME:	COMPLETE NAME:	COMPLETE NAME:
DATE:	DATE:	DATE:
SIGNATURE:	SIGNATURE:	SIGNATURE: