

Florfenicol 10% W/V

Oral Solution

(FOR VETERINARY USE ONLY)

COMPOSITION

Florfenicol 100 g/L
Benzyl Alcohol 8.5 ml

PRODUCT DESCRIPTION

The finished product is light yellowish colour liquid.

PHARMACODYNAMICS

Like chloramphenicol, florfenicol is a broad spectrum antibiotic that has activity against many bacteria. It acts by binding to the 50S ribosome, thereby inhibiting bacterial protein synthesis.

PHARMACOKINETICS

After the first oral gavage administration of florfenicol of 42mg/kg bw/day for 3 days to broiler chicken, the highest concentration of florfenicol in serum was measured after 30 minutes then declined to reach 810 µg/ml, 8 hours later. Florfenicol could not be detected in the serum, 60 hours after the last administration. During a continuous treatment of broiler chicken with florfenicol in drinking water at a daily dose of 26mg/kg bw for 18 hours for 3 days, the estimated mean serum concentration of florfenicol was 711µg/ml. No florfenicol was detected in the serum, 72 hours after the terminal dose. After repeated oral administration of ¹⁴C-florefnicol, given twice daily 12 hours apart at 20mg/kg bw/dose for three days, 93.7% and 98.2% of total radioactivity administered were excreted within 1 day and 7 days after the last dose. In excreta, at 7 days, the parent compound represented the major fraction of the radioactivity (42%), florfenicol amine (25%),

florfenicol oxamic acid (5%) and florfenicol alcohol (10%) and the remaining part of radioactivity being represented by a small percentage of three unknown compounds. No monochloroflorfenicol was detected in excreta. In a radiometric study, chicken received ¹⁴C-florefnicol by oral gavage twice daily 12 hour apart at 20 mg/kg bw/dose for three days (i.e. 40 mg/kg bw/day for three days). Twenty-four hour after the end of oral administration, the levels of radioactivity in edible tissues were 146 µg equivalents floefenicol/kg in muscle, 475µg/kg in skin+ fat, 11148 µg/kg in liver and 3125 µg/kg in kidney. Then declined to reach approximately 50 µg/kg in muscle and in skin + fat 5 days after the end of treatment, whereas in liver and kidney significant amounts of residues were still found : 2403 and 844µg equivalents florfenicol/kg respectively. Florfenicol amine was the major metabolite measured in edible tissues of chicken, florfenicol, the only microbiologically active compounds, being only detected in skin + fat in low concentrations (4%) and in kidney (1%). In a non-radiometric study, broiler chickens received florfenicol according to the recommended regimen via drinking water at concentrations equivalent to 17 to 30 mg/kg bw/day for 3 days. At 0.5 days after the end of the treatment the concentrations of florfenicol amine were lower than the limit of quantification for muscle (less than 50 µg/kg) and for skin + fat (less than 109 µg/kg) whereas in liver and kidney, significant amounts of florfenicol-amine were measured 2862 and 1161µg/kg respectively. Florfenicol amine concentrations declined to reach 2038 and 679 µg/kg in liver and kidney, 1 day after the end of the treatment and to 1215 and 484 µg/kg in liver and kidney, 3 days after the end of the treatment. Seven days after the end of the treatment, the concentrations of florfenicol amine were below the limit of quantification for liver (less than 461 µg/kg) and could be still measured in kidney (136 µg/kg).



SHENNONG ANIMAL HEALTH (M) SDN BHD (441647-X)

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1, 47610 Subang Jaya, Selangor, Malaysia.

INDICATIONS

Broiler chickens: Florfenicol 10% W/V Oral Solution is indicated in broiler chickens for the treatment and control of air sacculitis associated with *E. coli* susceptible to florfenicol.

CONTRA-INDICATIONS

Do not administer in cases of previous allergic reactions to florfenicol.

ADMINISTRATION & DOSAGE

1ml of Florfenicol 10% W/V Oral Solution in 1L water or 100ppm (100mg/L) for 3 consecutive days in drinking water. This dosage rate is equivalent to 35mg florfenicol/kg body weight of birds.

INTERACTION WITH OTHER MEDICAMENTS

No data available.

OVERDOSE AND TREATMENT

None known

PREGNANCY AND LACTATION

Studies in laboratory animals have not revealed any evidence of potential embryotoxic or foetotoxic effect of florfenicol. The use is not recommended during pregnancy and lactation. The product is not for use in animals from which eggs are produced for human consumption.

WITHDRAWAL TIME

Treated animals must not be slaughtered for use in food for 5 days for broiler chickens after the latest treatment with this drug.

SIDE EFFECT

None known

MAXIMUM RESIDUAL LIMIT (MRL)

Florfenicol MRL: Muscle (chicken)= 100 microgram/kg, kidney (chicken)= 750 microgram/kg, liver (chicken)= 2500 microgram/kg.

WARNING AND PRECAUTIONS

The medicated water should be administered as the only source of drinking water for 3 consecutive days. Inappropriate use of the veterinary medicinal product may increase the prevalence of bacteria resistant to florfenicol. To limit the development of antimicrobial resistance, the choice of this drug as the most appropriate treatment should be confirmed by clinical experience supported where possible by pathogen culture and drug susceptibility testing.

Not for human use

- Keep out of reach of children.

- Jauhi ubat dari kanak-kanak.

Avoid direct contact with skin, eyes and clothes. In case of accidental skin exposure, wash with soap and water. Remove contaminated clothing. Consult a physician if irritation persists.

SHELF LIFE

36 months from date of manufacture. Use within 3 months and keep tightly after first opening of container. Finish treatment within 12 hours after mixing or dilution.

STORAGE

Store in cool dry place (below 25°C). Strictly avoid light and exposure.

PACKAGING

Product is filled in plastic bottle of 1000 ml, sealed with aluminum foil and aluminum screw cap, fitted guarantee plastic cap.

DATE OF REVISION: 12 MAY 2019



SHENNONG ANIMAL HEALTH (M) SDN BHD (441647-X)

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10, Jalan Industri USJ 1/19, Taman Perindustrian USJ 1, 47610 Subang Jaya, Selangor, Malaysia.