

SALINOMYCIN (QILU) 12% GRANULES

Description and Composition

SALINOMYCIN (QILU) 12% GRANULES is a brownish free-flowing granules. Each kg contains 120g potency Salinomycin Sodium as its active ingredient.

Pharmacodynamics

Salinomycin sodium has a high affinity for cations of coccidium, especially for Na⁺, K⁺, Rb⁺, and cations outflow from the cells selectively, resulting in the interruption of concentration of cations.

Pharmacokinetics

A GLP-compliant study has been performed recently of which the aim was to revisit the metabolic fate of SAL in the chicken₃₈. Groups of 3 male and 3 female animals were administered by gavage twice daily and for 7 consecutive days [₁₄C]-SAL doses corresponding to a 70 mg SAL kg⁻¹ feed. No information was given on the labeling positions in the SAL molecule. The animals were slaughtered after 0, 1, 3, 5 and 7 days withdrawal of the additive. No significant sex difference was observed concerning the metabolic profiles in the excreta and tissues. The balance study confirmed indicating a rapid excretion (see below results from other studies that are inline with this information):

The balance study showed that 95% of the radioactivity was excreted within 48h of which 1.8% and 2.5% (results of two studies) was expired indicating a significant decarboxylation of the molecule.

A number of different non-GLP studies were performed. Tissue residues were the highest in the liver followed, in decreasing order, by the kidney, skin / fat and muscle.

The non-extractable radioactivity in the liver and kidney increased considerably along the withdrawal period to reach values of 51 - 76% and 65 - 70% after 48h, respectively, which indicates the formation of bound-residues or the incorporation of [₁₄C]-labelled fragments into normal tissue components.

Indication

As an aid in the prevention of coccidiosis in broiler chickens caused by *Eimeria necatrix*, *E. tenella*, *E. acervulina*, *E. brunetti*, *E. maxima* and *E. mivati*.

Recommended Dose

Prevention of coccidiosis : 500g per ton of feed

Route of Administration

For oral administration via feed.

Contraindication

Salinomycin must not be mixed with or used simultaneously with feed containing any other anticoccidial or substance having a similar effect. Salinomycin contains an ionophore. Simultaneous use with certain medicinal substances (eg tiamulin) maybe contra-indicated.

Warning and Precautions

For use in feeds for broiler chickens, chickens for fattening only. Dangerous for turkeys and equines.

Clean all mixing equipment thoroughly before preparing other feeding stuffs. Operators handling the product should wear protective clothing including facemask and gloves. Avoid inhalation and skin contact with the product. If accidental eye contact occurs, rinse out with water immediately. Do not leave or dispose of the product where it may be accessible to animals not under treatment or to wildlife.

Interactions with Other Medicaments

Salinomycin must not be mixed with or used simultaneously with feed containing any other anticoccidial or substance having a similar effect. Salinomycin is an ionophore. Simultaneous use with certain medicinal substances (eg tiamulin) maybe contra-indicated.

Pregnancy and Lactation

Do not feed to laying chickens.

Side Effects

N/A

Symptoms and Treatment of Overdose

N/A

Storage Condition

Store below 30°C in a dry place.

Shelf Life

2 years.

Withdrawal Period

Meat: Zero days

Packing

25kg

Manufacturer:

SHANDONG QILU KING-PHAR PHARMACEUTICAL CO., LTD.

No. 21 Qinglong Road, Pingyin country, Jinan, Shandong, P.R.C, (Postal Code): 250400

Product Registration Holder:

RANGE PHARMA SDN. BHD.

No. 1, Jalan TSB 11, Taman Industri Sg. Buloh, 47000 Sg. Buloh, Selangor Darul Ehsan, Malaysia.

Tel: 603-61568708 Fax: 603-61568707

Email: info@rangepharma.com

Revised date: 22.6.2026