

ANISOL 20% W.S.P

Soluble Powder

(FOR VETERINARY USE ONLY)

COMPOSITION

Levamisole (as hydrochloride) 200 mg/g

PRODUCT DESCRIPTION

Anisole W.S.P 200 is a white color soluble powder.

PHARMACODYNAMIC

Levamisole is an anthelmintic of the imidazothiazoles family; it is the levorotatory form of tetramisole. Because of its cholinomimetic mode of action by binding to the acetylcholine receptor, it acts at the nerve ganglia of the nematode resulting in paralysis caused the death of the parasite. Levamisole has no ovicidal activity.

PHARMACOKINETIC

After oral administration, gastrointestinal absorption of levamisole is rapid, peak plasma concentrations are reached in 2 to 4 hours. It undergoes metabolism in the liver and is rapidly excreted and mainly in the urine as metabolites.

INDICATIONS

Treatment of infections with the following parasites :

In cattle and sheep :

- Gastrointestinal roundworms : Trichostrongylus spp , Cooperia spp , Ostertagia ostertagi (unless inhibited larvae) Haemonchus spp , Nematodirus spp , Bunostomum spp , Oesophagostomum spp , Chabertia ovine (sheep) .
- Lungworms : Dictyocaulus viviparus .

In pigs:

- Gastrointestinal roundworms : Ascaris suum , Strongyloides ransomi , Oesophagostomum spp .
- Lungworms : Metastrongylus spp.

In poultry :

- Ascaridia spp , Heterakis spp , Capilaria spp , Amidostomum anseris (goose) .

CONTRA-INDICATIONS

Animals should not be treated simultaneously or within 14 days before or after use with organophosphorous compounds or diethylcarbamazine citrate.

ADMINISTRATION & DOSAGE

Cattle and sheep: 7.5mg of levamisole per kg bodyweight, in a single administration by oral route, without exceeding the total dose per animal of 3.75g in cattle and 0.375g in sheep, 0.37g of powder per 10kg bodyweight diluted in a little water without exceeding the dose of 18.75g in cattle and 1.87g in sheep.

Pigs: 7.5mg of levamisole per kg bodyweight, in a single administration by oral route, without exceeding the total dose per animal of 0.75g, 0.37g of powder per 10kg diluted in a little water or soup diluted without exceeding the dose of 3.75g per animal.

Poultry: 20mg of levamisole per kg bodyweight, approximately 1g of powder per liter of drinking water. The amount of medicated drinking water consumed by poultry depends on their physiological and clinical state. To obtain the recommended dose, the concentration of levamisole should be adjusted accordingly.

Use a spray dispenser clean and properly calibrated.



SHENNONG ANIMAL HEALTH (M) SDN BHD (441647-X)

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10 Jalan Industri USJ 1/19, Taman Perindustrian USJ 1, 47610 Subang Jaya, Selangor, Malaysia.

INTERACTION WITH OTHER MEDICAMENTS

The association of the product with anticholinesterase agents, such as external organophosphorus pesticides should be avoided.

STATEMENT ON USAGE DURING PREGNANCY AND LACTATION

No teratogenic effect was observed with levamisole in experimental animals (Mice, rats and rabbits). No embryotoxic or teratogenic have been reported in cattle, sheep and pigs. The veterinary medicinal product can be used in females during pregnancy and lactation.

OVERDOSE AND TREATMENT

At normal therapeutic dosages side effects are rarely seen. Overdose may occasionally result in the appearance of cholinergic-type symptoms such as salivation, muscular tremors and head shaking. They are more likely to be observed in cattle than in sheep.

SIDE EFFECT

Occasionally at the recommended dose cattle may show signs of lip-licking and slight muscle tremor.

WITHDRAWAL TIME

Meat and offal: 3 days

Milk: In the absence of a withdrawal period for the milk, do not use in females producing milk for consumption, lactating or in dry period nor future milk producing animals for consumption in the 2 months preceding the parturition.

Eggs: in the absence of a withdrawal period for the eggs, do not use in species producing eggs for consumption, within 4 weeks of onset of the laying and during it.

STORAGE

Store in room temperature ($\leq 25^{\circ}\text{C}$) and dry place.

SHELF LIFE

24 months from the date of manufacture. After first opening of container, keep the product tightly closed and in good storage conditions. Use up the remaining product content so long. And finish treatment through drinking water within 12 hours.

WARNING AND PRECAUTIONS

Special precautions for use in animals: Body weight must be assessed as accurately as possible before calculating the dose. Resistance can develop when used frequently and repeatedly. Special warning for each target species: Do not use in females producing milk. Do not use in layers. Special precautions to be taken by the person administering the veterinary medicinal product to animals: Avoid direct contact with skin. Wash hands after use. Personal protective equipment consisting of impermeable rubber gloves should be worn when handling the veterinary medicinal product. In case of accidental ingestion, seek medical advice immediately.

PACKAGING

The powder filled 250 g in aluminum sachets, by automatic filling machine and sealing, and packed in carton box.

DATE OF REVISION: 14 MAR 2019



SHENNONG ANIMAL HEALTH (M) SDN BHD (441647-X)

神農動物保健(馬)有限公司

No. 2, 4, 6, 8 & 10 Jalan Industri USJ 1/19, Taman Perindustrian USJ 1, 47610 Subang Jaya, Selangor, Malaysia.