

Pharmacode position may change as per Supplier's m/c requirement &additional
small pharma code may appear on the front / back panel



SUMMARY OF PRODUCT CHARACTERISTICS

XINCLAV 1.2g

Amoxicillin and Clavulanate potassium for Injection 1.2 g

Rx Only

NAME OF DRUG PRODUCT: Amoxicillin and Clavulanate potassium for injection 1.2 g

(TRADE) NAME OF PRODUCT: XINCLAV 1.2g

STRENGTH: 1.2 g.

PHARMACEUTICAL DOSAGE FORM: Injection

QUALITATIVE AND QUANTITATIVE COMPOSITIONS:

Amoxicillin and Clavulanate potassium for injection 1.2 g

Each vial contains:

Amoxicillin Sodium Ph.Eur. equivalent to Amoxicillin 1000 mg and Clavulanate Potassium Ph.Eur. equivalent to Clavulanic acid 200 mg.

PHARMACEUTICAL FORM:

A white to off-white crystalline moisture sensitive powder filled in 30 mL tubular vial, stoppered with 20 mm grey colour bromo butyl rubber stopper and sealed with 20 mm dark blue colour flip off seal.

Description of the reconstituted solution: A light yellow coloured clear solution.

Description of Diluent: Not Applicable, because Diluent will not come with the product.

CLINICAL PARTICULARS

Therapeutic Indications

Amoxicillin and clavulanate potassium is indicated for short-term treatment of bacterial infections at the following sites:

Upper respiratory tract infections (including ENT) e.g. recurrent tonsilitis, sinusitis, otitis media. Lower respiratory tract infections e.g. acute exacerbation of chronic bronchitis, lobar and bronchopneumonia.

Genito-urinary tract infections e.g. cystitis, urethritis, pyelonephritis. Skin and soft tissue infections e.g boils, abscesses, cellulitis, wound infections.

Bone and joint infections e.g osteomyelitis.

Other infections e.g intra-abdominal sepsis.

Amoxicillin and clavulanate potassium intravenous is also indicated for prophylaxis against infection which may be associated with major surgical procedures such as gastrointestinal, pelvic, head and neck, cardiac, renal, joint replacement and biliary tract.

Infections caused by amoxicillin-susceptible organisms are amenable to Amoxicillin and Clavulanate Potassium for Injection treatment due to its amoxicillin content. Mixed infections caused by amoxicillin-susceptible organisms in conjunction with Amoxicillin and Clavulanate Potassium for Injection-susceptible β -lactamase producing organisms may therefore be treated with Amoxicillin and Clavulanate Potassium for Injection.

Posology and method of administration

Dosages for the treatment of infection

Adults and children over 12 years: Usually 1.2 g eight hourly. In more serious infections, increase frequency to six-hourly intervals.

Children 3 months-12 years: Usually 30 mg/kg * Amoxicillin and Clavulanate potassium eight hourly. In more serious infections, increase frequency to six-hourly intervals.

Children 0-3 months: 30 mg/kg* Amoxicillin and Clavulanate potassium every 12 hours in premature infants and in full-term infants during the perinatal period, increasing to eight hours thereafter.

*Each 30 mg Amoxicillin and Clavulanate potassium provides 5 mg Clavulanic acid with 25 mg amoxicillin.

Adult dosage for surgical prophylaxis:

The usual dose is 1.2 g Amoxicillin and Clavulanate potassium Intravenous given at the induction of anaesthesia. Operations where there is a high risk of infection, e.g. colorectal surgery, may require three and up to four, doses of 1.2 g Amoxicillin and Clavulanate potassium Intravenous in a 24 hour period. These doses are usually given at 0, 8, 16 (and 24) hours. This regimen can be continued for several days if the procedure has a significantly increased risk of infection.

Clear clinical signs of infection at operation will require a normal course of intravenous or oral Amoxicillin and Clavulanate potassium therapy post-operatively.

Dosage in renal impairment:

Adults

Mild impairment (creatinine clearance >30 ml/min): No change in dosage.

Moderate impairment (creatinine clearance 10-30 ml/min): 1.2 g IV stat., followed by 600 mg IV 12 hourly.

Severe impairment (creatinine clearance <10 ml/min): 1.2 g IV stat., followed by 600 mg IV 24 hourly. Dialysis decreases serum concentrations of Amoxicillin and Clavulanate potassium and an additional 600 mg IV dose may need to be given during dialysis and at the end of dialysis.

Children

Similar reductions in dosage should be made for children.

Dosage in hepatic impairment

Dose with caution; monitor hepatic function at regular intervals.

There are, as yet, insufficient data on which to base a dosage recommendation.

Each 1.2 g vial of Amoxicillin and Clavulanate potassium contains 1.0 mmol of potassium and 2.7 mmol of sodium (approx).

Administration

Amoxicillin and Clavulanate potassium Intravenous may be administered either by intravenous injection or by intermittent infusion. It is not suitable for intramuscular administration.

Contraindications

Penicillin hypersensitivity. Attention should be paid to possible cross-sensitivity with other β -lactam antibiotics, e.g. cephalosporins.

A previous history of Amoxicillin and Clavulanate potassium or penicillin-associated jaundice/hepatic dysfunction.

Special warnings and precautions for use

Changes in liver function tests may occur in some patients receiving Amoxicillin and Clavulanate potassium. Amoxicillin and Clavulanate potassium should be used with caution in patients with evidence of hepatic dysfunction.

Cholestatic jaundice, which may be severe, but is usually reversible, may occurs rarely. Signs and symptoms may not become apparent for several weeks after treatment has ceased.

In patients with renal impairment, dosage should be adjusted according to the degree of impairment.

If the parenteral administration of high doses is necessary, the sodium content must be taken into account in patients on sodium restricted diet.

In patients with reduced urine output crystalluria may occurs very rarely, predominantly with parenteral therapy. During administration of high doses of amoxicillin it is advisable to maintain adequate fluid intake and urinary output in order to reduce the possibility of amoxicillin crystalluria. Amoxicillin precipitates in bladder catheters after intravenous administration of large doses. A regular check of patency should be maintained.

Serious and occasionally fatal hypersensitivity reactions (including anaphylactoid and severe cutaneous adverse reactions) have been reported in patients receiving therapy with beta-lactams. Before initiating therapy with Amoxicillin and Clavulanate potassium, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins, carbapenems or other beta-lactam agents. If an allergic reaction occurs, Amoxicillin and Clavulanate potassium must be discontinued immediately and appropriate alternative therapy instituted.

Erythematous rashes may be associated with glandular fever in patients receiving amoxicillin.

Prolonged use may also occasionally result in overgrowth of non-susceptible organisms.

Interaction with other medicinal products and other forms of interaction

Prolongation of bleeding time and prothrombin time occurs in some patients receiving Amoxicillin and Clavulanate potassium. Amoxicillin and Clavulanate potassium should be used with care in patients on anti-coagulation therapy.

In common with other broad-spectrum antibiotics, Amoxicillin and Clavulanate potassium may reduce the efficacy of oral contraceptives and patients should be warned accordingly.

Concomitant use of allopurinol during treatment with amoxicillin can increase the likelihood of allergic skin reactions.

Pregnancy and lactation

Prophylactic treatment with Amoxicillin and Clavulanate potassium may be associated with an increased risk of necrotising enterocolitis in neonates. As with all medicines, use should be avoided in pregnancy, especially during the first trimester, unless considered essential by the physician.

Amoxicillin and Clavulanate potassium may be administered during the period of lactation. With the exception of the risk of sensitisation, associated with the excretion of trace quantities in breast milk.

Effects on ability to drive and use machines

None known.

Undesirable effects

Side effects are uncommon and mainly of a mild and transitory nature.

A/s: 210 x 300 mm Black

 AUROBINDO Packaging Development		Product Name	Component	Item Code	Date & Time
		Xinclav 1.2	Leaflet	P1530394	17.02.2022 & 2:10 pm
		Customer / Country	Version No.	Reason Of Issue	Reviewed / Approved by
Malaysia_Unit 12	01	Revision			
Team Leader	Sravan	Dimensions	No. of Colours : 01		
Initiator	Saikanth	210 x 300 mm			
Artist: 	Pharmacode	 30394			
	30394				
Additional Information : Supersede item Code: P1522416					Sign / Date

Gastrointestinal reactions:

Diarrhoea, indigestion, nausea, vomiting and mucocutaneous candidiasis occur. Antibiotic-associated colitis (including pseudomembranous colitis and haemorrhagic colitis) may occur rarely.

Renal and urinary tract disorders:

Crystalluria occurs very rarely.

Hepatic effects:

Moderate and asymptomatic rises in AST and/or ALT and alkaline phosphatases occur occasionally. Hepatitis and cholestatic jaundice may occur rarely. These hepatic reactions are more common with Amoxicillin and Clavulanate potassium than with other penicillins.

Signs and symptoms usually occur during or shortly after treatment but in some cases may not occur until several weeks after treatment has ended. Hepatic reactions are usually reversible but they may be severe and very rarely, deaths occur.

Hypersensitivity reactions:

Urticarial and erythematous skin rashes sometimes occur. Rarely erythema multiforme, Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous exfoliative dermatitis, acute generalised exanthematous pustulosis (AGEP), serum sickness like syndrome and hypersensitivity vasculitis occur. Treatment should be discontinued if one of these disorders occurs. In common with other β -lactam antibiotics angioedema and anaphylaxis occurs. Interstitial nephritis can occur rarely.

Haematological effects:

As with other β -lactams transient leucopenia (including neutropenia and agranulocytosis), thrombocytopenia and haemolytic anaemia may occur rarely. Prolongation of bleeding time and prothrombin time may rarely occur.

CNS effects:

CNS effects occur very rarely. These include reversible hyperactivity, dizziness, headache and convulsions. Convulsions may occur with impaired renal function or in those receiving high doses.

Skin and subcutaneous tissue disorders:

Frequency 'very rare': Drug Reaction with Eosinophilia and Systemic Symptoms (DRESS)

Local: Thrombophlebitis at the site of injection occurs occasionally.

Over dosage

Gastrointestinal symptoms and disturbance of the fluid and electrolyte balances may occur. They may be treated symptomatically with attention to the water electrolyte balance. Amoxicillin and Clavulanate potassium may be removed from the circulation by haemodialysis.

PHARMACOLOGICAL PROPERTIES

Pharmacodynamic properties

Resistance to many antibiotics is caused by bacterial enzymes, which destroy the antibiotic before it can act on the pathogen. The Clavulanate in Amoxicillin and Clavulanate potassium anticipates this defence mechanism by blocking the β -lactamase enzymes, thus rendering the organisms sensitive to amoxicillin's rapid bactericidal effect at concentrations readily attainable in the body.

Clavulanate by itself has little antibacterial activity; however, in association with amoxicillin as Amoxicillin and Clavulanate potassium, it produces an antibiotic agent of broad spectrum with wide application in hospital and general practice.

Amoxicillin and Clavulanate potassium is bactericidal to a wide range of organisms including:

Gram-positive

Aerobes: *Enterococcus faecalis**, *Enterococcus faecium**, *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus viridans*, *Staphylococcus aureus**, Coagulase negative *staphylococci** (including *Staphylococcus epidermidis**), *Corynebacterium* species, *Bacillus anthracis**, *Listeria monocytogenes*.

Anaerobes: *Clostridium* species, *Peptococcus* species, *Peptostreptococcus*.

Gram-negative

Aerobes: *Haemophilus influenzae**, *Moraxella catarrhalis** (*Branhamella catarrhalis*), *Escherichia coli**, *Proteus mirabilis**, *Proteus vulgaris**, *Klebsiella* species*, *Salmonella* species*, *Shigella* species*, *Bordetella pertussis*, *Brucella* species, *Neisseria gonorrhoeae**, *Neisseria meningitidis**, *Vibrio cholerae*, *Pasteurella multocida*.

Anaerobes: *Bacteroides* species* including *B. fragilis*.

* Some members of these species of bacteria produce β -lactamase, rendering them insensitive to amoxicillin alone.

Pharmacokinetic properties

The pharmacokinetics of the two components of Amoxicillin and Clavulanate potassium is closely matched. Both Clavulanate and amoxicillin have low levels of serum binding; about 70% remains free in the serum.

Doubling the dosage of Amoxicillin and Clavulanate potassium approximately doubles the serum levels achieved.

PHARMACEUTICAL PARTICULARS

List of excipients

None.

Incompatibilities

Amoxicillin and Clavulanate potassium Intravenous should not be mixed with blood products, other proteinaceous fluids such as protein hydrolysates or with intravenous lipid emulsions.

If Amoxicillin and Clavulanate potassium is prescribed concurrently with an aminoglycoside, the antibiotics should not be mixed in the syringe, intravenous fluid container or giving set because loss of activity of the aminoglycoside can occur under these conditions.

Directions for Use:

Dissolve the contents in 20 ml Sterile Water for Injection.

Amoxicillin and Clavulanate Potassium for injection should be given by slow intravenous injection over a period of 3 to 4 minutes.

After reconstitution, immediately use the freshly prepared solution.

Shelf life

24 months.

Special precautions for storage

Store in a dry place below 30°C.

After reconstitution, immediately use the freshly prepared solution.

Nature and contents of container

Mono vial and 10 vials without Sterile Water for Injection.

Manufactured by


AUROBINDO
Aurobindo Pharma Ltd,
Unit XII, Survey No. 314,
Bachupally, Bachupally Mandal,
Medchal-Malkajgiri District,
Telangana State, India.

Regd. Office: Plot No.: 2, Maitrivihar,
Ameerpet, Hyderabad-500 038,
Telangana State, India.

Product registration holder:
Healol Pharmaceuticals Sdn Bhd, 74-3,
Jalan Wangsa Delima 6,
KLSC Wangsa Maju,
53300 Kuala Lumpur

Date of revision: February 2022