

PACKAGE INSERT

Maduramicin Ammonium Premix 1% Powder

Brand or product Name	Maduramicin Ammonium Premix 1% Powder
Name and strength of active substance(s)	Maduramicin ammonium 1%
Product Description	Pale yellow to russet brown powder or granule
Pharmacodynamics	Maduramicin is a monovalent glycoside polyether ionophore antibiotic. It induces an anticoccidial effect on the asexual forms (sporozoites, merozoites and falciform bodies) of the endogenous development of coccidia in poultry, some resistant forms to other ionophore antibiotics included. Its anticoccidial activity is due to disorder in ion metabolism (potassium and sodium in particular).
Pharmacokinetics	Maduramicin was excreted rapidly in chickens. Steady state was observed after three days in the excreta, but after six days in plasma. Maduramicin was the main compound excreted (26 %). Odemethylation represented the main metabolic pathway of maduramicin common to chickens and rats. Liver is the target tissue; maduramicin is the marker residue in all tissues tested. Tissue residue kinetics of total residues and marker residue indicate a rapid decline of residues in all tissues. Ratios maduramicin to total residues have been established.
Environmental Properties	Not Applicable
Indication	As an aid in the prevention of coccidiosis caused by <i>Eimeria necatrix</i> , <i>E. tenella</i> , <i>E. acervulina</i> , <i>E. brunetti</i> , <i>E. maxima</i> and <i>E. mivati</i> .
Recommended Dosage	Thoroughly mix the product into the feed at a level of 500 gm per tonne of broiler feed to obtain a concentration of 5 g/t maduramicin ammonium in the complete feed. Feed containing the product should be the only ration fed to broilers to avoid dilution of the active ingredient.
Mode of Administration	Oral route, by mixing the product with feeding stuff
Contraindication	None.

Warning and precaution	For use in feeds intended for chicken only. Do not mix with or use simultaneously with feed containing another anticoccidial. Direct contact with the skin is to be avoided. When mixing and handling this product, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling this product. If accidental eye contact occurs, rinse thoroughly with water. Do not allow horses or other equines access to feeds containing this product. Ingestion of maduramicin ammonium may be fatal.
Interaction with other medicaments	None.
Statement on usage during pregnancy and lactation	Not applicable.
Adverse effects/Undesirable Effects	Not known.
Overdose and treatment	None.
Incompatibilities (for injections only)	N/A
Withdrawal Period(s)	Withdraw 5 day before slaughter.
MRL	Maduramicin: Edible tissue, muscle (chicken) 240ug/kg Fat (chicken) 480ug/kg Liver (chicken) 720ug/kg
Storage conditions (may be omitted if the information is stated on the label or outer carton labels)/(should be supported by stability data)	Store at temperature 25°C.
Dosage form and packaging available	Dosage form: Granules/powder in the form of powder Packaging size: 25kg
Name and Address of manufacturer	Zhejiang Apelo Kangyu Bio-pharmaceutical Co., Ltd. Geshan Industrial Zone, Dongyang, Zhejiang 322109, China
Name and address registration holder	Weissen Company (M) Sdn Bhd Lot 105, No. 84, Jalan Velox 23, Taman Velox, 48000 Rawang, Selangor, Malaysia

Date of revision of package insert	01/07/2024
Shelf life of the veterinary product as packaged for sale	2 years
Shelf life after first opening the container(where relevant)	Not Applicable
Shelf life after dilution or reconstitution according the directions (where relevant)	Not Applicable