

ZENTA 100KIU SOLUBLE POWDER

FOR ANIMAL USE ONLY

Brand or Product Name :

ZENTA 100KIU SOLUBLE POWDER

Name and Strength of Active Substance(s)

Each gram of ZENTA100KIU SOLUBLE POWDER contains gentamicin sulphate 100,000 IU (which is equivalent to 14.1% w/w gentamicin)

Product Description

ZENTA 100KIU Soluble Powder is a white or almost white powder.

Pharmacodynamics properties

Gentamicin sulfate is an aminoglycoside obtained from cultures of *Micromonospora purpurea*. Gentamicin acts on susceptible bacteria by irreversibly binding to the 30S ribosomal subunit thereby inhibiting protein synthesis. It is considered to be a bactericidal antibiotic. Gentamicin is active against most aerobic Gram-negative bacteria, some aerobic Gram-positive bacteria and against mycoplasma.

Pharmacokinetics properties

Gentamicin, like other aminoglycosides, is not appreciably absorbed after oral administration.

Indication

Swine (weanling) : For control and treatment of colibacillosis caused by strains of *E. coli* sensitive to gentamicin

Swine : For control and treatment of swine dysentery associated with *Treponema hyodysenteriae*

Recommended Dosage

Swine (weanling) : 0.05 g ZENTA 100 KIU SOLUBLE POWDER per Liter of drinking water (equivalent to 6.6 mg gentamicin / Liter) for 3 consecutive days, to provide 1.1 mg gentamicin per kg of body weight per day.

Swine : 0.1 g ZENTA 100 KIU SOLUBLE POWDER per Liter of drinking water (equivalent to 13.2 mg gentamicin / Liter) for 3 consecutive days, to provide 2.2 mg gentamicin per kg of body weight per day.

Administration

To be administered orally via animal drinking water. Do not store or offer medicated drinking water in rusty containers since the drug is quickly destroyed in such containers.

Medicated drinking water should be prepared daily and be the sole source of drinking water for 3 consecutive days. Treatment may be repeated if dysentery recurs.

Mixing Directions : To assure thorough dissolution, place the ZENTA 100KIU Soluble Powder in a small mixing container and add 25 liter of water . Stir until all the powder completely dissolved. Mix this concentrated stock solution with water to make the final volume of treated animal drinking water. Medicated water should be made up immediately prior to provision. Medication of the water supply should be continuous during the treatment period and no other source of water should be available. The product should be given over a minimum period of 6 hours per day to allow all animals to drink.

Contraindications

The use of this drug is contraindicated in animals with a history of an allergic reaction to aminoglycosides.

Interaction With Other Medicinal Product

Concurrent use with other nephrotoxic and ototoxic drugs may increase risk of nephrotoxicity or ototoxicity.

Pregnancy and Lactation

Do not use in pregnant animals.

Warnings and Precautions

Zenta 100KIU Soluble Powder should be used with caution in animals with preexisting renal disease. Not to be handled by person with known aminoglycosides allergy. Use appropriate personal protection. Wear impervious gloves when handling powder. Open sachet with care and not in windy area to avoid forming dust cloud. Use adequate masks and safety goggles in case of powder being blown.

Respiratory Protection	:	Wear dust mask in case powder being blown.
Hands	:	Wear Impervious gloves when handling powder.
Eye Protection	:	Wear Safety goggles when handling powder.
Skin Protection	:	Wear protective clothing to minimize skin exposure.

Side Effects

There have been no reported side effects following the recommended use of this product in swine.

Overdose and Treatment

None known.

Withdrawal Period(s)

10 days

Shelf-life

Shelf life as packaged for sale: 3 years from the date of production.

Shelf life after first opening of the immediate packaging: 24 hours

Shelf life after reconstitution in drinking water: 24 hours

Any opened sachet or unused medicated water to be discarded after 24 hours.

Maximum Residual Limit (MRL)

Pig : Muscle, fat = 100µg/kg ,kidney = 1000µg/kg, liver = 200µg/kg [Ref. REGOV, Version 2, Dec. 2009]

Storage Conditions

Store in a dry place below 30°C. Keep out of reach of children. Jauhkan ubat daripada kanak-kanak.

Disposal of Containers

Traces of medicated powder left in an opened aluminum sachet can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned aluminum sachet can be safely disposed in trash according to local government regulations

Dosage Forms and Packaging Available

Available in 100g, 200g, 500g and 1000g powder per sachet, pack in 10, 20, 50 and 100 sachets per box.

Name and Address of Manufacturer

Life Biopharma Sdn. Bhd. (Company No. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan Darul Khusus, Malaysia.

Date of Revision of Package Insert

2014-04-14