

[FRONT PAGE](#)

148 mm

75 mm

Pharmacode
Read Direction →

90208521

- Do not use on kittens aged below 8 weeks.
- Do not use in animals with existing impairment of cartilage growth.
- Pregnant and nursing animals should not be treated.

- Item code placement is flexible and can be moved to suit artwork

Template:	148 x 210 mm	Packaging Spec(s):			TPM_016192
Barcodes/Type:	Laetus: 0016	Add. Info:			N/A
		Minimum Core Data Point Size:			8 pt
GTIN:	N/A	Proof #	V1	By/Date	ORT / MK / 19.07.2022
BLACK NON-PRINTING INFO					

EL ANGO ARTWORK | LEGEND v13

148 mm

- Do not use on animals suffering from central seizure disorders as enrofloxacin can stimulate the central nervous system.
- Enrofloxacin is partially excreted via the kidneys; as with all fluoroquinolones, excretion may therefore be delayed in animals with existing renal damage.
- Do not apply to animals intended for human consumption.
- Animals known to be hypersensitive to fluoroquinolones should not be treated with enrofloxacin.
- Do not use in the event of known or suspected resistance to quinolones, as there is almost complete cross-resistance to other fluoroquinolones in animals resistant to quinolones.
- Do not administer concomitantly with tetracyclines, phenicols or macrolides as there is a risk of antagonistic effects.
- Do not administer prophylactically.

Warning and Precautions

Special precautions for use:

Observe the official and local antibiotics guidelines when using the veterinary medicinal product. Using this drug product in a way that deviates from the conditions stipulated in the SPC can promote the prevalence of fluoroquinolone-resistant bacteria and may reduce the efficacy of treatment with other quinolones because of potential cross-resistance. Pyoderma is generally secondary to an underlying primary infection. It is advisable to identify the primary underlying cause and treat the animal accordingly.

Special precautions for use in animals:

Baytril flavour 50 mg tablets should only be used after prior confirmation of the diagnosis by bacteriological sensitivity testing of the causal organisms and in the presence of resistance to other antibiotics.

Like all fluoroquinolones, Baytril flavour 50 mg tablets should not be used in trivial infections because of the potential for resistance development.

Caution is required when administered the veterinary drug product to animals with severe liver or kidney damage.

Special precautions to be taken by the person administering the veterinary medicinal product to animals:

Individuals with known hypersensitivity to fluoroquinolones or any of the other ingredients should avoid contact with this veterinary drug. Do not eat, drink or smoke during administration. Wash hands after use.

In the event of accidental ingestion, consult a doctor immediately and show him this pack insert.

Avoid contact with the eyes. In the event of contact with the eyes, rinse the affected eye immediately with plenty of water.

Interaction with Other Medicaments

- The elimination of theophylline may be delayed.
- Combinations of Baytril (enrofloxacin) with chloramphenicol, macrolide antibiotics or tetracyclines may produce antagonistic effects.
- Simultaneous administration of substances containing magnesium or aluminium may impair the absorption of enrofloxacin.
- There is complete cross-resistance between fluoroquinolones.
- Concomitant administration of fluoroquinolones with non-steroidal anti-inflammatory drugs (NSAIDs), e.g. flunixin, to dogs is subject to strict veterinary supervision. Due to interactions between the two substances, the elimination half-lives can be extended which can lead to undesired drug reactions.
- Concomitant administration of fluoroquinolones with digoxin should be avoided due to the resulting risk of an elevated oral bioavailability of digoxin.

Pregnancy and Lactation

Pregnant and nursing animals should not be treated.

Adverse reactions

Gastrointestinal disorders may occur in isolated cases.

Overdose and Treatment

In the event of large overdoses, the initial symptoms are likely to be loss of appetite and vomiting. In very rare cases, diarrhoea or CNS symptoms (dilated pupils, muscle tremor, impaired coordination and convulsions) can occur after treatment, making it necessary to stop treatment. Administration of antacids containing magnesium or aluminium is recommended to decrease absorption of enrofloxacin after oral ingestion.

Withdrawal Period

Not applicable.

Storage Condition

Do not store above 30°C.

Dosage Forms and packaging available

Baytril™ Flavour Tablets 50 mg is available in pack of 50 tablets per box.

Marketing Authorisation Holder:

Elanco Malaysia Sdn Bhd

Unit 5.04, Level 5 & 6, Tower Block, The Bousteador, No. 10, Jalan PJU 7/6, Mutiara Damansara, 47800 Petaling Jaya, Selangor, Malaysia

Manufactured by:

KVP Pharma + Veterinär Produkte GmbH

Projensdorfer Straße 324, 24106 Kiel, Germany.

Date of revision: 12 March 2021

Baytril is sold by Elanco or its affiliates and is not a product of Bayer. The Product Name Baytril is owned by Bayer and used under license.

Elanco™

PRODUCT INFO

BLUE #:	502171APK	Product Name:	Baytril flavour 50 mg
Item Code:	90208521	Component:	Leaflet
Product Code:	LBAH	Pack Size:	N/A
Previous Item Code	90204238		

Page 2 of 2

ARTWORK INFO

Template:	148 x 210 mm	Packaging Spec(s):	TPM_016192
Barcodes/Type:	Laetus: 0016	Add. Info:	N/A
		Minimum Core Data Point Size:	8 pt
GTIN:	N/A	Proof #	V1
		By/Date	ORT / MK / 19.07.2022
 BLACK NON-PRINTING INFO			

ELANCO ARTWORK LEGEND v13