

ToltraCox-25 Oral Solution

FOR ANIMAL USE ONLY

Brand or Product Name :

ToltraCox-25 Oral Solution

Name and Strength of Active Substance(s)

Each ml contains 25 mg toltrazuril.

Product Description

ToltraCox-25 Oral Solution is a clear, colorless to light yellowish or brownish liquid.

Pharmacodynamics

Toltrazuril induces changes in the fine structure of coccidial development stages that are mainly due to the swelling of the endoplasmic reticulum and of the Golgi apparatus and to abnormalities in the perinuclear space and disturbances in nuclear division. Toltrazuril also causes a reduction of enzymes in the respiratory chain of the parasites.

Pharmacodynamic studies in laboratory animals indicated that toltrazuril had no anticoagulant, fibrinolytic, analgesic, diuretic or anticonvulsant effect. It had no effect on gastrointestinal transit time. Oral doses of up to 30 mg/kg bw had no haemodynamic or cardiac effects in anaesthetised dogs although 100mg/kg bw caused a slight tensor effect as reflected by an increase in peripheral resistance.

Pharmacokinetics

In poultry toltrazuril is absorbed at a rate of at least 50%. The highest equivalent concentrations are found in liver. The active substance is rapidly metabolised in poultry. The main metabolite is characterised as a sulfone-derivative. About 1 week after the last dose this metabolite represents by far the most relevant residue in the animal.

Indication

For the treatment of Coccidiosis in Broilers and Broiler Breeders.

It is effective against *Eimeria acervulina*, *E. brunetti*, *E. maxima*, *E. necatrix*, *E. tenella*, *E. mitis*.

Recommended Dosage

The recommended dose rate is 7 mg Toltrazuril per kg bodyweight per day given for 2 consecutive days. This corresponds to : 28 ml ToltraCox-25 Oral Solution (equivalent to 700 mg toltrazuril) per 100 kg of bodyweight per day for 2 consecutive days.

As a guide for usage this is normally equivalent to 25 ppm (equivalent to 1 ml ToltraCox-25 Oral Solution) per litre of drinking water for continuous medication over 48 hours, or to 75 ppm (equivalent to 3 ml ToltraCox-25 Oral Solution) per litre of drinking water given for 8 hours per day, on 2 consecutive days.

Treatment Regimen

Curative treatment

Chicken : 1 medication period (over 2 days).

Only make up sufficient drinking water for requirements.

After calculation of the required amount of product, this volume should be carefully dispensed from the container into a domestic measuring jug reserved and labelled specifically for the purpose. Measurement must be made carefully to ensure correct dosage. Product must not be stored in the jug.

Mode of Administration

To be administered via animal drinking water. Mixing direction : ToltraCox-25 must be added in the amount of animal drinking water estimated to be consumed within 12 – 24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water must be prepared each day.

Contraindications

Do not use in birds producing eggs for human consumption.

Warnings and Precautions

Special warnings for each target species

The product does not interfere with the development of immunity against coccidiosis. As with all anticoccidials, prolonged use may result in the development of resistant strains.

Special precautions for use

i. Special precautions for use in animals

Dilutions more concentrated than 1:1,000 (1 ml ToltraCox-25 Oral Solution to 1 litre drinking water) may result in precipitation. Predilution is not recommended.

ii. Special precautions to be taken by the person administering the medicinal product to animals

Wear synthetic rubber gloves when handling the product. Wash any splashes from skin or eyes immediately with water. Wash hands and exposed skin after use.

Do not eat, drink or smoke whilst using the product.

iii. Other precautions

Only make up sufficient drinking water for requirements. Wash out empty container with water 3 times when medicating birds and add washings to drinking water.

Store litter from treated houses as long as possible. However, it should only be stored where there is no risk of polluting surface water or groundwater. Do not apply to agricultural land at a rate greater than 1.5 tonnes/hectare/year. The principle metabolite of toltrazuril is persistent and mobile in groundwater.

Interactions with Other Medicaments

None known.

Statement on usage during pregnancy and lactation

Not applicable. Product is for use in chickens (not birds producing eggs for human consumption).

Side Effects

None known.

Symptoms and Treatment of Overdose

A three - five fold overdosage is readily tolerated without any symptoms. If the recommended dose is exceeded beyond 3 - 5 times there is a decrease in water intake.

Withdrawal Period(s)

Meat: 18 days. Do not use in birds producing eggs for human consumption.

Shelf-life

2 years from date of manufacture. Opened bottle must be discarded after 24 hours. Medicated water must be prepared fresh daily. Any unused medicated water is to be discarded after 24 hours.

Allowable Maximum Residual Limit (MRL)

Chicken : Muscle = 100 µg /kg, Skin+fat = 200 µg /kg, Liver = 600 µg /kg, Kidney = 400 µg /kg. [Ref. EMEA/MRL/743/00-FINAL, dated June 2000]

Storage Conditions

Store in below 25°C. Keep medicine away from children. Jauhkan ubat dari kanak-kanak.

Disposal of containers

Traces of medicated liquid left in an opened bottle can be rinsed out with clean animal drinking water and use in animal treatment. Cleaned empty bottle can be safely disposed in trash according to local government regulations.

Packaging available

Available in 1 litre per bottle, pack in 12 bottles or 20 bottles per carton. Also available in 5 litre per bottle, pack in 4 bottle per carton.

Name and Address of manufacturer

Life Biopharma Sdn. Bhd. (company no. 546163-D)

No. 250-251, Jalan Industri Galla 13, Kawasan Perindustrian Galla, 70200 Seremban, Negeri Sembilan, Malaysia.

Date of Revision of Package Insert

2013-11-11