

Product Name:

NEOMIVET 50% Water Soluble powder

Composition:

Each gram contains:

Neomycin (as sulfate) 500mg

Product Description:

White to light Yellow powder

Pharmacodynamics:

Neomycin has a mechanism of action and spectrum of activity (primarily gram-negative aerobes) similar to the other aminoglycosides, but in comparison to either gentamicin or amikacin, it is significantly less effective against several species of gram-negative organisms, including strains of *Klebsiella*, *E. coli*, and *Pseudomonas*. However, most strains of neomycin-resistant bacteria of these species remain susceptible to amikacin.

Pharmacokinetics:

Approximately 3% of a dose of neomycin is absorbed after oral or rectal (retention enema) administration, but this can be increased if gut motility is slowed or if the bowel wall is damaged. Therapeutic levels are not attained in the systemic circulation after oral administration. After IM administration, therapeutic levels can be attained with peak levels occurring within 1 hour of dosing. The drug apparently distributes to tissues and is eliminated like the other aminoglycosides (refer to Amikacin monograph for more details). Orally administered neomycin is nearly all excreted unchanged in the feces.

Indications:

Cattle (excluding veal calves), Sheep, Swine and Goats:

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days.

Turkeys: 10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) for 5 days.

Dosage and Administration:

Cattle (exclude veal calves), Swine, Sheep and Goats: 10 milligrams of neomycin sulfate per pound of body weight per day (22 milligrams per kilogram) in divided doses for a maximum of 14 days.

Contraindications:

Do not use in animals with known hypersensitivity to Neomycin.

Interaction with Other Medicaments:

None known.

Pregnancy and Lactation:

None known.

Side effects:

None known.

Symptoms and Treatment of Overdose:

No known symptoms of overdose.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

Withdraw all milk and milk replacer from scouring animals. Use of the product should be based on susceptibility testing of the bacteria isolated from the animal. If this is not possible, therapy should be based on local epidemiological information about susceptibility of the target bacteria.

Turkeys: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 5 consecutive days.

Cattle (excluding veal calves), Swine, Sheep and Goats: Add to drinking water or milk. Not for use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after use in liquid supplements. Prepare a fresh solution daily. If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian, but not to exceed a total of 14 days.

Container disposal: Dispose of any unused product and empty container in accordance with local waste regulation authority.

Withdrawal Periods:

Cattle excluding veal calves: 1 day before slaughter

Sheep: 2 days before slaughter

Swine and goats: 3 days before slaughter

Meat: Do not use less than 5 days before slaughter for human consumption.

Turkeys: zero days before slaughter

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,

Prai Industrial Estate MK1,

13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

NAM PHARMA SDN. BHD.

5, Lebuhr Perusahaan Klebang II,

Taman Perindustrian Antarabangsa IGB

31200 Chemor, Perak.

DATE OF REVISION: APRIL 2019