

**Product Name:**

TILMIVET 50% Water Soluble powder

**Composition:**

Each gram contains:

Tilmicosin (as phosphate) 500mg

**Product Description:**

Yellow to reddish powder

**Pharmacodynamics:**

NA

**Pharmacokinetics:**

When administered orally to chickens or turkeys and pigs via the drinking water at recommended concentrations, tilmicosin is absorbed and moves rapidly out of the serum into areas of low pH. This results in very low serum concentrations, but detectable levels are found in lung tissue as early as 6 hours after commencing treatment. In chickens or turkeys, tilmicosin is also detected in pooled airsac tissue as early as 6 hours after commencing treatment. It is also known that tilmicosin is concentrated in alveolar macrophages. Tilmicosin is also found at fairly high concentrations in liver and kidney tissue, since it is excreted both via the bile into the faeces and also via the urine.

**Indications:**

Pigs Prevention and treatment of respiratory disease caused by *Actinobacillus pleuropneumoniae*, *Mycoplasma hyopneumoniae*, *Pasteurella multocida* and other organisms sensitive to tilmicosin

**Dosage and Administration:**

Pigs: To be included in the drinking water to provide a daily dose of 15-20 mg/kg bodyweight for 5 days, which may be achieved by the inclusion of 200 mg tilmicosin per litre.

**Contraindications:**

Do not use in horses.

**Interaction with Other Medicaments:**

None known.

**Pregnancy and Lactation:**

None known.

**Side effects:**

None known.

**Symptoms and Treatment of Overdose:**

No known symptoms of overdose.

**Warning and Precautions:**

Keep out of reach of children.

Jauhi dari kanak-kanak.

Tilmicosin should not be administered by injection to pigs. Whenever possible, tilmicosin should only be used based on susceptibility testing. May be highly toxic when administered parenterally and, therefore, should not be injected. Do not allow horses and other equines access to drinking water containing tilmicosin. Tilmicosin may induce irritation. Macrolides, such as tilmicosin, may also cause hypersensitivity (allergy) following injection, inhalation, ingestion or contact with skin or eye. Hypersensitivity to tilmicosin may lead to cross reactions to other macrolides and vice versa. Allergic reactions to these substances may occasionally be serious and therefore direct contact should be avoided. To avoid exposure during preparation of the medicated drinking water, wear overalls, safety glasses, and impervious gloves. Do not eat, drink or smoke when handling this product. Wash hands after use. In the case of accidental ingestion, wash out mouth immediately with water and seek medical advice. In the event of accidental skin contact, wash thoroughly with soap and water. In case of accidental eye contact, flush the eyes with plenty of clean, running water. Do not handle the product if you are allergic to ingredients in the product. If you develop symptoms following exposure, such as skin rash, you should seek medical advice and show the physician this warning. Swelling of the face, lips and eyes or difficulty in breathing are more serious symptoms and require urgent medical attention. Manure from treated animals should not be deposited on the same field in successive years.

**Withdrawal Periods:**

Pigs: 21 days

**Storage conditions:**

Store in cool, dark place. Store below 30° C. Protect from direct light.

**Pack Size:**

1kg, 100gm

**Distributed by Product Registration Holder:**

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,

Prai Industrial Estate MK1,

13600 Prai, Pulau Pinang, Malaysia.

**Manufactured By:**

THYE PHARMA SDN. BHD.

No. 2, Lorong Industri Ringan Permatang Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam, Pulau Pinang.

DATE OF REVISION: APRIL 2019