

Product Name:

TIAMIVET 50% Water Soluble powder

Composition:

Each gram contains:

Tiamulin 500mg

Product Description:

White powder

Pharmacodynamics:

Tiamulin hydrogen fumarate is a semi-synthetic diterpene antibiotic. The mode of action is by inhibition of ribosomal protein synthesis in sensitive bacteria. It is a bacteriostatic antibiotic and the following organisms show sensitivity in vitro:

Brachyspira: Brachyspira hyodysenteriae, B. pilosicoli

Mycoplasmas: Mycoplasma hyopneumoniae, M. hyorhinis, M. hyosynoviae, Ureaplasma spp., M. gallisepticum, M. synoviae, and M. meleagridis.

Gram-positive: Staphylococcus spp., Streptococcus spp., Arcanobacterium pyogenes.

Gram-negative: Pasteurella spp., Klebsiella pneumoniae, Actinobacillus (Haemophilus) spp., Fusobacterium necrophorum, Bacteroides spp., Campylobacter coli, Lawsonia intracellularis.

Pharmacokinetics:

Following oral administration, tiamulin hydrogen fumarate is rapidly absorbed and peak serum concentrations are achieved 2 - 4 hours after administration and concentrates in some tissues e.g. lung. Following administration, it is rapidly metabolised and excreted.

Indications:

Pigs: For the treatment, prevention and control of swine dysentery caused by Brachyspira hyodysenteriae and complicated by Fusobacterium and Bacteroides spp. Chickens: For the reduction in the severity of disease caused by mycoplasmas. Turkeys: For the reduction in the severity of disease caused by mycoplasmas.

Dosage and Administration:

Pigs:

The dosage is 8.8 mg of the active substance per kg bodyweight daily, administered in the drinking water of pigs for 3 to 5 days, depending on the severity of the infection and/or the duration of the disease. If there is no response to treatment within 5 days, the diagnosis should be re-established.

Chickens:

The dosage for chickens is 25 mg of active substance per kg bodyweight. Tiamulin should be administered for 3-5 days and will provide approximately the following daily dosage of tiamulin depending on the age of the bird.

Day old chick: 125-150 mg/kg bodyweight

4 week old broiler: 30-50 mg/kg bodyweight

10 week old pullet: 30-45 mg/kg bodyweight

Layer: 25 mg/kg bodyweight

Turkeys:

The dosage for turkeys is 25 mg of active substance per kg bodyweight. Tiamulin should be administered for 5 days and will provide approximately the following daily dosage of tiamulin depending on the age of the bird.

1 week old poult: 70 mg/kg bodyweight

4 week old poult: 50 mg/kg bodyweight

8 week old poult: 20-30 mg/kg bodyweight

20 week old poult: 25 mg/kg bodyweight

Monitor water intake at frequent intervals during medication.

Contraindications:

Not to be used in animals with known sensitivity to the active ingredient.

Animals and birds should not receive products containing monensin, narasin or salinomycin, during or for at least seven days before or after treatment. Severe growth depression or death may result.

Interaction with Other Medicaments:

Animals and birds should not receive products containing monensin, narasin or salinomycin during or for at least seven days before or after treatment. Severe growth depression or death may result. Concomitant use of tiamulin and the ionophore anticoccidial maduramicin may lead to a mild to moderate growth depression in chickens. The situation is transient and recovery normally occurs within 3-5 days following withdrawal of tiamulin treatment. This does not seem to occur with the ionophores lasalocid or semduramicin.

Pregnancy and Lactation:

The product can be used in pregnant and lactating pigs.

Tiamulin may be used in laying and breeding birds as it has been shown to have no adverse effects on egg production, fertility and hatchability in chickens and turkeys.

Side effects:

On rare occasions erythema or mild oedema of the skin may occur in pigs following the use of tiamulin hydrogen fumarate. Water intake may be depressed during the administration of tiamulin to birds. It appears to be concentration dependant with 0.025% tiamulin reducing intake by approximately 15%. It does not appear to have any adverse effect on overall performance of the birds or efficacy of the product.

Symptoms and Treatment of Overdose:

There is a relatively high therapeutic index with tiamulin and the likelihood of an overdosage is considered remote, especially as water intake and hence tiamulin intake is reduced if abnormally high concentrations are given. The LD₅₀ for chickens is 1290 mg/kg bodyweight and for turkeys 840 mg/kg bodyweight.

A single oral dose of 100 mg/kg BW caused hyperpnoea and abdominal complaints in pigs. At a dose of 150 mg/kg the only effects on the central nerve system was lethargy. A dose of 55 mg/kg during 14 days caused increased salivation and a mild irritation of the stomach. Tiamulin hydrogen fumarate has a relatively high therapeutic index in pigs. The minimum lethal dose has not been established in pigs.

The clinical signs of toxicity in chickens are: vocalization, clonic cramps and lying in a lateral position. Symptoms in turkeys are: clonic cramps, lateral or dorsal position, salivation and ptosis.

If signs of intoxication do occur, promptly remove the medicated water and replace with fresh water.

Warning and Precautions:

Keep out of reach of children.

Jauhi dari kanak-kanak.

In order to avoid interactions between tiamulin and the incompatible ionophores monensin, narasin and salinomycin, the feed mill supplying the feed should be notified that tiamulin will be used and that these products should not be included in the feed or contaminate the feed.

The feed should be tested for the ionophores prior to use if there is any suspicion that contamination of the feed might occur. If an interaction does occur, stop tiamulin water medication immediately and replace with fresh water. Remove contaminated feed as soon as possible and replace with feed not containing the tiamulin-incompatible ionophores.

When mixing, direct contact with the skin and eyes should be avoided by wearing impermeable rubber gloves and safety glasses.

In case of accidental eye contact, irrigate the eyes thoroughly with clean running water immediately. Seek medical advice if irritation persists.

Contaminated clothing should be removed and any splashes on to the skin should be washed off immediately.

Wash hands after use.

Withdrawal Periods:

Animal and birds must not be slaughtered for human consumption during treatment.

Pigs 7 Chicken: May be slaughtered for human consumption only after 2 days from the last treatment.

Turkeys: may be slaughtered for human consumption only after 5 days from the last treatment.

Eggs: Zero days

Storage conditions:

Store in cool, dark place. Store below 30° C. Protect from direct light.

Pack Size:

1kg, 100gm

Distributed by Product Registration Holder:

LIM SENG PHARMACHEM SDN. BHD.

2692, Tingkat Perusahaan Enam,

Prai Industrial Estate MK1,

13600 Prai, Pulau Pinang, Malaysia.

Manufactured By:

THYE PHARMA SDN. BHD.

No. 2, Lorong Industri Ringan Permatang Tinggi 8,

Kawasan Industri Ringan Permatang Tinggi,

14000 Bukit Mertajam, Pulau Pinang.

DATE OF REVISION: APRIL 2019